


**Image** 






## MED12 Antibody

| CSB-PA936493  |
|---|
| Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.   |
| Q93074  |
| Synthetic peptide of Human MED12  |
| Rabbit  |
| Human, Mouse  |
| ELISA,IHC;ELISA:1:2000-1:5000,IHC:1:50-1:200  |
| The initiation of transcription is controlled in part by a large protein assembly known as the preinitiation complex. A component of this preinitiation complex is a 1.2 MDa protein aggregate called Mediator. This Mediator component binds with a CDK8 subcomplex which contains the protein encoded by this gene, mediator complex subunit 12 (MED12), along with MED13, CDK8 kinase, and cyclin C. |
| Liquid  |
| Non-conjugated  |
| -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol  |
| Antigen affinity purification   |
| IgG   |
| Homo sapiens (Human)  |
| MED12   |
|   |



The image on the left is immunohistochemistry of paraffin-embedded Human brain tissue using CSB-PA936493(MED12 Antibody) at dilution 1/40, on the right is treated with synthetic peptide. (Original magnification: x200)



The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using CSB-PA936493(MED12 Antibody) at dilution 1/40, on the right is treated with synthetic peptide. (Original magnification: ×200)