





FADD (Ab-191) Antibody

Product Code CSB-PA975869 Storage Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. Uniprot No. Q61160 Immunogen Synthesized peptide derived from C-terminal of Human FADD. Raised In Rabbit Species Reactivity Human, Mouse Specificity The antibody detects endogenous levels of total FADD protein. Tested Applications ELISA,WB;WB:1:500-1:3000 Relevance Apoptotic adaptor molecule that recruits caspase-8 or caspase-10 to the activated Fas (CD95) or TNFR-1 receptors. The resulting aggregate called the death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation. Active caspase-8 initiates the subsequent cascade of caspases mediating apoptosis By similarity. Involved in interferon-mediated antiviral immune response, playing a role in the positive regulation of interferon signaling By similarity. Involved in interferon-mediated antiviral immune response, playing a role in the positive regulation of interferon signaling By similarity. Involved in interferon mediated antiviral immune response, playing a role in the positive regulation of interferon signaling By similarity. Science 309:1559-1563(2005) Form Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Purification Method The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Clonality Polyclona		
Uniprot No. Q61160 Immunogen Synthesized peptide derived from C-terminal of Human FADD. Raised In Rabbit Species Reactivity Human,Mouse Specificity The antibody detects endogenous levels of total FADD protein. Tested Applications ELISA,WB;WB:1:500-1:3000 Relevance Apoptotic adaptor molecule that recruits caspase-8 or caspase-10 to the activated Fas (CD95) or TNFR-1 receptors. The resulting aggregate called the death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation. Active caspase-8 initiates the subsequent cascade of caspases mediating apoptosis By similarity. Involved in interferon-mediated antiviral immune response, playing a role in the positive regulation of interferon signaling By similarity. Zhang J., Mol. Cell. Biol. 16:2756-2763(1996). Hsu H., Cell 84:299-308(1996). Carninci P., Science 309:1559-1563(2005) Form Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Purification Method The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Clonality Polyclonal Alias FADD; Fas (TNFRSF6)-associated via death domain; FAS-associated death domain protein; FAS-associating death domain-containing protein; Fas-associating protein with death domain Product Type Polyclonal Antibody	Product Code	CSB-PA975869
Immunogen Synthesized peptide derived from C-terminal of Human FADD. Raised In Rabbit Species Reactivity Human,Mouse Specificity The antibody detects endogenous levels of total FADD protein. Tested Applications ELISA,WB;WB:1:500-1:3000 Relevance Apoptotic adaptor molecule that recruits caspase-8 or caspase-10 to the activated Fas (CD95) or TNFR-1 receptors. The resulting aggregate called the death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation. Active caspase-8 initiates the subsequent cascade of caspases mediating apoptosis By similarity. Involved in interferon-mediated antiviral immune response, playing a role in the positive regulation of interferon signaling By similarity. Zhang J., Mol. Cell. Biol. 16:2756-2763(1996). Hsu H., Cell 84:299-308(1996). Carminci P., Science 309:1559-1563(2005) Form Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Purification Method The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Clonality Polyclonal Alias FADD; Fas (TNFRSF6)-associated via death domain; FAS-associated death domain protein; FAS-associating death domain-containing protein; Fas-associating protein with death domain Product Type Polyclonal Antibody Species Mus musculus (Mouse) <tr< th=""><th>Storage</th><td>Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.</td></tr<>	Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Raised In Rabbit Species Reactivity Human,Mouse Specificity The antibody detects endogenous levels of total FADD protein. Tested Applications ELISA,WB;WB:1:500-1:3000 Relevance Apoptotic adaptor molecule that recruits caspase-8 or caspase-10 to the activated Fas (CD95) or TNFR-1 receptors. The resulting aggregate called the death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation. Active caspase-8 initiates the subsequent cascade of caspases mediating apoptosis By similarity. Involved in interferon-mediated antiviral immune response, playing a role in the positive regulation of interferon signaling By similarity. Zhang J., Mol. Cell. Biol. 16:2756-2763(1996). Hsu H., Cell 84:299-308(1996). Carninci P., Science 309:1559-1563(2005) Form Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCI, 0.02% sodium azide and 50% glycerol. Purification Method The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Clonality Polyclonal Alias FADD; Fas (TNFRSF6)-associated via death domain; FAS-associated death domain protein; FAS-associating death domain-containing protein; Fas-associating protein with death domain Product Type Polyclonal Antibody Species Mus musculus (Mouse) Target Names Fadd/FADD	Uniprot No.	Q61160
Specificity Human,Mouse Specificity The antibody detects endogenous levels of total FADD protein. Tested Applications ELISA,WB;WB:1:500-1:3000 Relevance Apoptotic adaptor molecule that recruits caspase-8 or caspase-10 to the activated Fas (CD95) or TNFR-1 receptors. The resulting aggregate called the death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation. Active caspase-8 initiates the subsequent cascade of caspases mediating apoptosis By similarity. Involved in interferon-mediated antiviral immune response, playing a role in the positive regulation of interferon signaling By similarity. Form Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Purification Method The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Clonality Polyclonal Alias FADD; Fas (TNFRSF6)-associated via death domain; FAS-associated death domain protein; FAS-associating death domain-containing protein; Fas-associating protein with death domain Product Type Polyclonal Antibody Species Mus musculus (Mouse) Target Names Fadd/FADD	Immunogen	Synthesized peptide derived from C-terminal of Human FADD.
Specificity The antibody detects endogenous levels of total FADD protein. Tested Applications ELISA,WB;WB:1:500-1:3000 Relevance Apoptotic adaptor molecule that recruits caspase-8 or caspase-10 to the activated Fas (CD95) or TNFR-1 receptors. The resulting aggregate called the death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation. Active caspase-8 initiates the subsequent cascade of caspases mediating apoptosis By similarity. Involved in interferon-mediated antiviral immune response, playing a role in the positive regulation of interferon signaling By similarity. Form Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Purification Method The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Clonality Polyclonal Alias FADD; Fas (TNFRSF6)-associated via death domain; FAS-associated death domain protein; FAS-associating death domain-containing protein; Fas-associating protein with death domain Product Type Polyclonal Antibody Species Mus musculus (Mouse) Target Names Fadd/FADD	Raised In	Rabbit
Tested Applications ELISA,WB;WB:1:500-1:3000 Apoptotic adaptor molecule that recruits caspase-8 or caspase-10 to the activated Fas (CD95) or TNFR-1 receptors. The resulting aggregate called the death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation. Active caspase-8 initiates the subsequent cascade of caspases mediating apoptosis By similarity. Involved in interferon-mediated antiviral immune response, playing a role in the positive regulation of interferon signaling By similarity. Zhang J., Mol. Cell. Biol. 16:2756-2763(1996). Hsu H., Cell 84:299-308(1996). Carninci P., Science 309:1559-1563(2005) Form Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Purification Method The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Clonality Polyclonal Alias FADD; Fas (TNFRSF6)-associated via death domain; FAS-associated death domain protein; FAS-associating death domain-containing protein; Fas-associating protein with death domain Product Type Polyclonal Antibody Species Mus musculus (Mouse) Fadd/FADD	Species Reactivity	Human, Mouse
Relevance Apoptotic adaptor molecule that recruits caspase-8 or caspase-10 to the activated Fas (CD95) or TNFR-1 receptors. The resulting aggregate called the death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation. Active caspase-8 initiates the subsequent cascade of caspases mediating apoptosis By similarity. Involved in interferon-mediated antiviral immune response, playing a role in the positive regulation of interferon signaling By similarity. Zhang J., Mol. Cell. Biol. 16:2756-2763(1996). Hsu H., Cell 84:299-308(1996). Carninci P., Science 309:1559-1563(2005) Form Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Purification Method The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Clonality Polyclonal Alias FADD; Fas (TNFRSF6)-associated via death domain; FAS-associated death domain protein; FAS-associating death domain-containing protein; Fas-associating protein with death domain Product Type Polyclonal Antibody Species Mus musculus (Mouse) Target Names Fadd/FADD	Specificity	The antibody detects endogenous levels of total FADD protein.
activated Fas (CD95) or TNFR-1 receptors. The resulting aggregate called the death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation. Active caspase-8 initiates the subsequent cascade of caspases mediating apoptosis By similarity. Involved in interferon-mediated antiviral immune response, playing a role in the positive regulation of interferon signaling By similarity. Zhang J., Mol. Cell. Biol. 16:2756-2763(1996). Hsu H., Cell 84:299-308(1996). Carninci P., Science 309:1559-1563(2005) Form Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Purification Method The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Clonality Polyclonal Alias FADD; Fas (TNFRSF6)-associated via death domain; FAS-associated death domain protein; FAS-associating death domain-containing protein; Fas-associating protein with death domain Product Type Polyclonal Antibody Species Mus musculus (Mouse) Target Names Fadd/FADD	Tested Applications	ELISA,WB;WB:1:500-1:3000
Purification Method The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Clonality Polyclonal Alias FADD; Fas (TNFRSF6)-associated via death domain; FAS-associated death domain protein; FAS-associating death domain-containing protein; Fas-associating protein with death domain Product Type Polyclonal Antibody Species Mus musculus (Mouse) Target Names Fadd/FADD	Relevance	activated Fas (CD95) or TNFR-1 receptors. The resulting aggregate called the death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation. Active caspase-8 initiates the subsequent cascade of caspases mediating apoptosis By similarity. Involved in interferon-mediated antiviral immune response, playing a role in the positive regulation of interferon signaling
chromatography using epitope-specific immunogen. Clonality Polyclonal Alias FADD; Fas (TNFRSF6)-associated via death domain; FAS-associated death domain protein; FAS-associating death domain-containing protein; Fas-associating protein with death domain Product Type Polyclonal Antibody Species Mus musculus (Mouse) Target Names Fadd/FADD		Hsu H., Cell 84:299-308(1996).
Alias FADD; Fas (TNFRSF6)-associated via death domain; FAS-associated death domain protein; FAS-associating death domain-containing protein; Fas-associating protein with death domain Product Type Polyclonal Antibody Species Mus musculus (Mouse) Target Names Fadd/FADD	Form	Hsu H., Cell 84:299-308(1996). Carninci P., Science 309:1559-1563(2005) Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4,
domain protein; FAS-associating death domain-containing protein; Fas-associating protein with death domain Product Type Polyclonal Antibody Species Mus musculus (Mouse) Target Names Fadd/FADD		Hsu H., Cell 84:299-308(1996). Carninci P., Science 309:1559-1563(2005) Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. The antibody was affinity-purified from rabbit antiserum by affinity-
Species Mus musculus (Mouse) Target Names Fadd/FADD	Purification Method	Hsu H., Cell 84:299-308(1996). Carninci P., Science 309:1559-1563(2005) Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Target Names Fadd/FADD	Purification Method Clonality	Hsu H., Cell 84:299-308(1996). Carninci P., Science 309:1559-1563(2005) Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Polyclonal FADD; Fas (TNFRSF6)-associated via death domain; FAS-associated death domain protein; FAS-associating death domain-containing protein; Fas-
	Purification Method Clonality Alias	Hsu H., Cell 84:299-308(1996). Carninci P., Science 309:1559-1563(2005) Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Polyclonal FADD; Fas (TNFRSF6)-associated via death domain; FAS-associated death domain protein; FAS-associating death domain-containing protein; Fas-associating protein with death domain
Image	Purification Method Clonality Alias Product Type	Hsu H., Cell 84:299-308(1996). Carninci P., Science 309:1559-1563(2005) Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Polyclonal FADD; Fas (TNFRSF6)-associated via death domain; FAS-associated death domain protein; FAS-associating death domain-containing protein; Fas-associating protein with death domain Polyclonal Antibody
	Purification Method Clonality Alias Product Type Species	Hsu H., Cell 84:299-308(1996). Carninci P., Science 309:1559-1563(2005) Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Polyclonal FADD; Fas (TNFRSF6)-associated via death domain; FAS-associated death domain protein; FAS-associating death domain-containing protein; Fas-associating protein with death domain Polyclonal Antibody Mus musculus (Mouse)




CUSABIO TECHNOLOGY LLC









Western blot analysis of extracts from A549 cells (Lane 2), using FADD (Ab-191) antiobdy. The lane on the left is treated with synthesized peptide.