

Image








PLCG1 (Ab-783) Antibody

Product Code	CSB-PA995465
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P19174
Immunogen	Peptide sequence around aa. 781~785 (G-F-Y-V-E) derived from Human PLCg1.
Raised In	Rabbit
Species Reactivity	Human,Mouse,Rat
Specificity	The antibody detects endogenous level of total PLCg1 protein.
Tested Applications	ELISA,WB;WB:1:500-1:1000
Relevance	PLC-gamma is a major substrate for heparin-binding growth factor 1 (acidic fibroblast growth factor)-activated tyrosine kinase.
	D D 11/15
	DeBell KE, et al. Mol Cell Biol. 1999 Nov; 19(11): 7388-7398. Ver
Form	
Form Purification Method	Ver Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+),
	Ver Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-
Purification Method	Ver Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific peptide.
Purification Method Clonality	Ver Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific peptide. Polyclonal
Purification Method Clonality Alias	Ver Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific peptide. Polyclonal Phosphoinositide phospholipase C; Phospholipase C-gamma-1;
Purification Method Clonality Alias Product Type	Ver Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific peptide. Polyclonal Phosphoinositide phospholipase C; Phospholipase C-gamma-1; Polyclonal Antibody



Western blot analysis of extracts from Hela and 293 cells using PLCg1(Ab-783) Antibody.