




# Recombinant Human Receptor-interacting serine/threonine-protein kinase 3 (RIPK3)

|                          |   |
|--------------------------|---|
| <b>Product Code</b>      | CSB-BP897497HU  |
| <b>Relevance</b>         | Essential for necroptosis, a programmed cell death process in response to death-inducing TNF-alpha family members. Upon induction of necrosis, RIPK3 interacts with, and phosphorylates RIPK1 and MLKL to form a necrosis-inducing complex. RIPK3 binds to and enhances the activity of three metabolic enzymes: GLUL, GLUD1, and PYGL. These metabolic enzymes may eventually stimulate the tricarboxylic acid cycle and oxidative phosphorylation, which could result in enhanced ROS production.   |
| <b>Abbreviation</b>      | RIPK3   |
| <b>Storage</b>           | The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.   |
| <b>Uniprot No.</b>       | Q9Y572  |
| <b>Alias</b>             | RIP-like protein kinase 3 Receptor-interacting protein 3  |
| <b>Product Type</b>      | Recombinant Protein   |
| <b>Immunogen Species</b> | Homo sapiens (Human)  |
| <b>Purity</b>            | Greater than 90% as determined by SDS-PAGE.   |
| <b>Sequence</b>          | MSCVKLWPSGAPAPLVSIEELENQELVGKGGFGTVFRAQHRKWGYDVAVKIV<br>NSKAISREVKAMASLDNEFVLRLEGVIEKVNWDQDPKPALVTKFMENGSLSGL<br>LQSQCPRPWPLLCRLLKEVVLGMFYLDQNPVLLHRDLKPSNVLLDPELHVKL<br>ADFGGLSTFQGGSSQSGTSGGEPGGTLGYLAPELFVNVNRKASTASDVYSFGIL<br>MWAVLAGREVELPTEPSLVYEAVCNRQNRPSLAELPQAGPETPGLEGLKELM<br>QLCWSSEPKDRPSFQECLPKTDEVFQMVENNMMNAAVSTVKDFLSQLRSSNR<br>RFSIPESGQGGTEMDGFRRTIENQHRSRNDVMVSEWLNKLNLEPPSSVPPKKC<br>PSLTKRSRAQEEQVPQAWTAGTSSDSMAQPPQTPETSTFRNQMPSPSTSTGT<br>PSPGPRGNQGAERQGMNWSCRTPENPVTGRPLVNIYNCSGVQVGDNNYLT<br>MQQTALPTWGLAPSGKGRGLQHPPPVGSQEGPKDPEAWSRPQGWNHSGK |
| <b>Lead Time</b>         | Delivery time may differ from different purchasing way or location, please kindly consult your local distributors for specific delivery time.   |
| <b>Research Area</b>     | Cell Biology  |
| <b>Source</b>            | Baculovirus   |
| <b>Gene Names</b>        | RIPK3   |
| <b>Protein Names</b>     | Recommended name: Receptor-interacting serine/threonine-protein kinase 3<br>EC= 2.7.11.1Alternative name(s): RIP-like protein kinase 3 Receptor-interacting protein 3 Short name= RIP-3   |



|                            |   |
|----------------------------|---|
| <b>Expression Region</b>   | 1-518aa   |
| <b>Notes</b>               | Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week. |
| <b>Tag Info</b>            | N-terminal 10xHis-tagged and C-terminal Myc-tagged  |
| <b>Mol. Weight</b>         | 60.9kDa   |
| <b>Protein Description</b> | Full Length   |

**Image**


(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

**Reconstitution**

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.