



Human intercellular adhesion molecule 1, ICAM-1 ELISA Kit

Product Code	CSB-E04574h
Abbreviation	ICAM1
Protein Biological Process 1	Cell Adhesion
Uniprot No.	P05362
Product Type	ELISA Kit
Immunogen Species	Homo sapiens (Human)
Sample Types	serum, plasma, tissue homogenates
Detection Range	0.78 ng/mL-50 ng/mL
Sensitivity	0.195 ng/mL
Assay Time	1-5h
Sample Volume	50-100ul
Detection Wavelength	450 nm
Lead Time	3-5 working days after you place the order, and it takes another 3-5 days for delivery via DHL or FedEx.
Research Area	Immunology
Quality Control	<p>A microplate reader capable of measuring absorbance at 450 nm, with the correction wavelength set at 540 nm or 570 nm.</p> <p>An incubator can provide stable incubation conditions up to 37°C±5°C.</p> <p>Centrifuge</p> <p>Vortex</p> <p>Squirt bottle, manifold dispenser, or automated microplate washer</p> <p>Absorbent paper for blotting the microtiter plate</p> <p>50-300ul multi-channel micropipette</p> <p>Pipette tips</p> <p>Single-channel micropipette with different ranges</p> <p>100ml and 500ml graduated cylinders</p> <p>Deionized or distilled water</p> <p>Timer</p> <p>Test tubes for dilution</p>
Gene Names	ICAM1
Tag Info	quantitative
Protein Description	Sandwich
Component	A micro ELISA plate ---The 96-well plate has been pre-coated with an anti-human ICAM1 antibody. This dismountable microplate can be divided into 12 x 8 strip plates.



Two vials lyophilized standard ---Dilute a bottle of the standard at dilution series, read the OD values, and then draw a standard curve.

One vial Biotin-labeled ICAM1 antibody (100 x concentrate) (120 µl/bottle) ---Act as the detection antibody.

One vial HRP-avidin (100 x concentrate) (120 µl/bottle) ---Bind to the detection antibody and react with the TMB substrate to make the solution chromogenic.

One vial Biotin-antibody Diluent (15 ml/bottle) ---Dilute the Biotin-antibody.

One vial HRP-avidin Diluent (15 ml/bottle) ---Dilute the HRP-avidin solution.

One vial Sample Diluent (50 ml/bottle)---Dilute the sample to an appropriate concentration.

One vial Wash Buffer (25 x concentrate) (20 ml/bottle) ---Wash away unbound or free substances.

One vial TMB Substrate (10 ml/bottle) ---Act as the chromogenic agent. TMB interacts with HRP, eliciting the solution turns blue.

One vial Stop Solution (10 ml/bottle) ---Stop the color reaction. The solution color immediately turns from blue to yellow.

Four Adhesive Strips (For 96 wells) --- Cover the microplate when incubation.
An instruction manual

Description

This Human ICAM1 ELISA Kit was designed for the quantitative measurement of Human ICAM1 protein in serum, plasma, tissue homogenates. It is a Sandwich ELISA kit, its detection range is 0.78 ng/mL-50 ng/mL and the sensitivity is 0.195 ng/mL.

Msds

{"0":{"fileurl":"https://www.cusabio.com/uploadfile/msds/MSDS CSB-E04574h.pdf","filename":"MSDS"}}