





# Pig interleukin 2,IL-2 ELISA Kit

Product Code	CSB-E06783p
Abbreviation	IL2
Protein Biological Process 1	Immunity
Target Name	interleukin 2
Uniprot No.	P26891
Alias	IL-2, TCGF, lymphokine, T cell growth factor aldesleukin interleukin-2 involved in regulation of T-cell clonal expansion
<b>Product Type</b>	ELISA Kit
Immunogen Species	Sus scrofa (Pig)
Protein Biological Process 3	Adaptive immunity
Sample Types	serum, plasma, tissue homogenates
<b>Detection Range</b>	15.6 pg/mL-1000 pg/mL
Sensitivity	3.9 pg/mL
Assay Time	1-5h
Sample Volume	50-100ul
<b>Detection Wavelength</b>	450 nm
Lead Time	3-5 working days after you place the order, and it takes another 3-5 days for delivery via DHL or FedEx.
Research Area	Immunology
Gene Names	IL2
Tag Info	quantitative
<b>Protein Description</b>	Sandwich
Description	This Pig IL2 ELISA Kit was designed for the quantitative measurement of Pig IL2 protein in serum, plasma, tissue homogenates. It is a Sandwich ELISA kit, its detection range is 15.6 pg/mL-1000 pg/mL and the sensitivity is 3.9 pg/mL.
Target Details	This protein is a secreted cytokine that is important for the proliferation of T and B lymphocytes. The receptor of this cytokine is a heterotrimeric protein complex whose gamma chain is also shared by interleukin 4 (IL4) and interleukin 7 (IL7). The expression of this gene in mature thymocytes is monoallelic, which represents an unusual regulatory mode for controlling the precise expression of a single gene. The targeted disruption of a similar gene in mice leads to ulcerative colitis-like disease, which suggests an essential role of this gene in

#### **CUSABIO TECHNOLOGY LLC**









the immune i	response to	antigenic	stimuli.
--------------	-------------	-----------	----------

#### **Product Precision**

Intra-assay Precision (Precision within an assay): CV%<8%

Three samples of known concentration were tested twenty times on one plate to assess.

Inter-assay Precision (Precision between assays): CV%<10%

Three samples of known concentration were tested in twenty assays to

assess.

## Linearity

To assess the linearity of the assay, samples were spiked with high concentrations of pig IL-2 in various matrices and diluted with the Sample Diluent to produce samples with values within the dynamic range of the assay.

?	Sample	Serum(n=4)
1:1	Average %	92
	Range %	80-99
1:2	Average %	86
	Range %	82-91
1:4	Average %	91
	Range %	86-97
1:8	Average %	94
	Range %	88-98

## Recovery

The recovery of pig IL-2 spiked to levels throughout the range of the assay in various matrices was evaluated. Samples were diluted prior to assay as directed in the Sample Preparation section.

Sample Type	Average % Recovery	Range
Serum (n=5)	94	89-102
EDTA plasma (n=4)	96	91-102


## **Typical**

These standard curves are provided for demonstration only. A standard curve should be generated for each set of samples assayed.









pg/ml OD1 OD2 Average Corrected

1000 2.884 2.915 2.900 2.764 500 2.123 2.104 2.114 1.978 250 1.249 1.272 1.261 1.125 125 0.818 0.846 0.832 0.696 62.5 0.492 0.440 0.466 0.330 31.2 0.314 0.322 0.318 0.182 15.6 0.263 0.258 0.261 0.125 ? 0 0.135 0.136 0.136

**Msds** 

{"0":{"fileurl":"https://www.cusabio.com/uploadfile/msds/MSDS CSB-E06783p.pdf","filename":"MSDS"}}