





# Pig interleukin 4,IL-4 ELISA Kit

Product Code	CSB-E06785p
Abbreviation	IL4
Protein Biological Process 1	Immunity
Target Name	interleukin 4
Uniprot No.	Q04745
Alias	BCGF-1, BCGF1, BSF1, IL-4, MGC79402, B cell growth factor 1 B_cell stimulatory factor 1 lymphocyte stimulatory factor 1
Product Type	ELISA Kit
Immunogen Species	Sus scrofa (Pig)
Protein Biological Process 3	B-cell activation
Sample Types	serum, plasma, cell culture supernates, tissue homogenates
<b>Detection Range</b>	6.25 pg/mL-400 pg/mL
Sensitivity	1.56 pg/mL
Assay Time	1-5h
Sample Volume	50-100ul
<b>Detection Wavelength</b>	450 nm
Lead Time	3-5 working days after you place the order, and it takes another 3-5 days for delivery via DHL or FedEx.
Research Area	Immunology
Gene Names	IL4
Tag Info	quantitative
<b>Protein Description</b>	Sandwich
Description	This Pig IL4 ELISA Kit was designed for the quantitative measurement of Pig IL4 protein in serum, plasma, cell culture supernates, tissue homogenates. It is a Sandwich ELISA kit, its detection range is 6.25 pg/mL-400 pg/mL and the sensitivity is 1.56 pg/mL .
Target Details	This protein is a pleiotropic cytokine produced by activated T cells. This cytokine is a ligand for interleukin 4 receptor. The interleukin 4 receptor also binds to IL13, which may contribute to many overlapping functions of this cytokine and

IL13. STAT6, a signal transducer and activator of transcription, has been shown to play a central role in mediating the immune regulatory signal of this cytokine.

chromosome 5q, with this gene particularly close to IL13. This gene, IL13 and

This gene, IL3, IL5, IL13, and CSF2 form a cytokine gene cluster on

#### **CUSABIO TECHNOLOGY LLC**









IL5 are found to be regulated coordinately by several long-range regulatory elements in an over 120 kilobase range on the chromosome. Two alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported.

#### **Product Precision**

Intra-assay Precision (Precision within an assay): CV%<8%

Three samples of known concentration were tested twenty times on one plate to assess.

Inter-assay Precision (Precision between assays): CV%<10%

Three samples of known concentration were tested in twenty assays to assess.

# Linearity

To assess the linearity of the assay, samples were spiked with high concentrations of pig IL-4 in various matrices and diluted with the Sample Diluent to produce samples with values within the dynamic range of the assay.

?	Sample	Serum(n=4)
1:1	Average %	86
	Range %	82-91
1:2	Average %	90
	Range %	85-95
1:4	Average %	89
	Range %	84-93
1:8	Average %	96
	Range %	92-101

### Recovery

The recovery of pig IL-4 spiked to levels throughout the range of the assay in various matrices was evaluated. Samples were diluted prior to assay as directed in the Sample Preparation section.

Sample Type	Average % Recovery	Range
Serum (n=5)	100	95-105
EDTA plasma (n=4)	92	86-96

## **Typical**

These standard curves are provided for demonstration only. A standard curve should be generated for each set of samples assayed.






🕜 Tel: +1-301-363-4651 🛛 Email: cusabio@cusabio.com 🕒 Website: www.cusabio.com 🥑







pg/ml OD1 OD2 Average Corrected

2.719 2.652 2.686 2.492 200 1.962 1.861 1.912 1.718 100 1.213 1.204 1.209 1.015 50 0.748 0.689 0.719 0.525 25 0.551 0.527 0.539 0.345 12.5 0.355 0.374 0.365 0.171 6.25 0.295 0.272 0.284 0.090 0 0.192 0.196 0.194 ?

Msds

{"0":{"fileurl":"https://www.cusabio.com/uploadfile/msds/MSDS CSB-E06785p.pdf","filename":"MSDS"}}