



# Rabbit basic fibroblast growth factor,bFGF ELISA Kit

<b>Product Code</b>	CSB-E06938Rb
<b>Abbreviation</b>	FGF2
<b>Protein Biological Process 1</b>	Angiogenesis
<b>Target Name</b>	fibroblast growth factor 2 (basic)
<b>Uniprot No.</b>	P48799
<b>Alias</b>	BFGF, FGFB, HBGF-2, basic fibroblast growth factor bFGF fibroblast growth factor 2 heparin-binding growth factor 2 prostatropin
<b>Product Type</b>	ELISA Kit
<b>Immunogen Species</b>	Oryctolagus cuniculus (Rabbit)
<b>Protein Biological Process 3</b>	Angiogenesis
<b>Sample Types</b>	serum, plasma, tissue homogenates
<b>Detection Range</b>	7.8 pg/mL-500 pg/mL
<b>Sensitivity</b>	1.95 pg/mL
<b>Assay Time</b>	1-5h
<b>Sample Volume</b>	50-100ul
<b>Detection Wavelength</b>	450 nm
<b>Lead Time</b>	3-5 working days after you place the order, and it takes another 3-5 days for delivery via DHL or FedEx.
<b>Research Area</b>	Cardiovascular
<b>Gene Names</b>	FGF2
<b>Tag Info</b>	quantitative
<b>Protein Description</b>	Sandwich

## Description

This Rabbit FGF2 ELISA Kit was designed for the quantitative measurement of Rabbit FGF2 protein in serum, plasma, tissue homogenates. It is a Sandwich ELISA kit, its detection range is 7.8 pg/mL-500 pg/mL and the sensitivity is 1.95 pg/mL.

## Target Details

This protein is a member of the fibroblast growth factor (FGF) family. FGF family members bind heparin and possess broad mitogenic and angiogenic activities. This protein has been implicated in diverse biological processes, such as limb and nervous system development, wound healing, and tumor growth. The



mRNA for this gene contains multiple polyadenylation sites, and is alternatively translated from non-AUG (CUG) and AUG initiation codons, resulting in five different isoforms with distinct properties. The CUG-initiated isoforms are localized in the nucleus and are responsible for the intracrine effect, whereas, the AUG-initiated form is mostly cytosolic and is responsible for the paracrine and autocrine effects of this FGF.

### Product Precision

Intra-assay Precision (Precision within an assay): CV%<8%

Three samples of known concentration were tested twenty times on one plate to assess.

Inter-assay Precision (Precision between assays): CV%<10%

Three samples of known concentration were tested in twenty assays to assess.

### Linearity

To assess the linearity of the assay, samples were spiked with high concentrations of rabbit bFGF in various matrices and diluted with the Sample Diluent to produce samples with values within the dynamic range of the assay.

	Sample	Serum(n=4)
1:1	Average %	87
	Range %	84-90
1:2	Average %	104
	Range %	100-108
1:4	Average %	85
	Range %	81-89
1:8	Average %	99
	Range %	95-102


### Recovery

The recovery of rabbit bFGF spiked to levels throughout the range of the assay in various matrices was evaluated. Samples were diluted prior to assay as directed in the Sample Preparation section.

Sample Type	Average % Recovery	Range
Serum (n=5)	83	80-86
EDTA plasma (n=4)	97	94-100

### Typical

These standard curves are provided for demonstration only. A standard curve should be generated for each set of samples assayed.



pg/ml	OD1	OD2	Average	Corrected
500	1.863	1.812	1.838	1.679
250	1.699	1.598	1.649	1.490
125	1.428	1.436	1.432	1.273
62.5	1.115	1.168	1.142	0.983
31.2	0.804	0.831	0.818	0.659
15.6	0.563	0.574	0.569	0.410
7.8	0.362	0.372	0.367	0.208
0	0.159	0.158	0.159	

## Msds

```
{
  "0": {
    "fileurl": "https://www.cusabio.com/uploadfile/msds/MSDS CSB-E06938Rb.pdf",
    "filename": "MSDS"
  }
}
```