





## Mouse Neprilysin(MME) ELISA kit

Product Code	CSB-EL014653MO
Abbreviation	MME
Target Name	membrane metallo-endopeptidase
Uniprot No.	Q61391
Alias	CALLA, CD10, DKFZp686O16152, MGC126681, MGC126707, NEP, atriopeptidase common acute lymphocytic leukemia antigen enkephalinase membrane metallo-endopeptidase (neutral endopeptidase, enkephalinase) m
Product Type	ELISA Kit
Immunogen Species	Mus musculus (Mouse)
Sample Types	serum, plasma,tissue homogenates
<b>Detection Range</b>	0.156 ng/mL-10 ng/mL
Sensitivity	0.039 ng/mL
Assay Time	1-5h
Sample Volume	50-100ul
<b>Detection Wavelength</b>	450 nm
Lead Time	3-5 working days after you place the order, and it takes another 3-5 days for delivery via DHL or FedEx.
Research Area	Others
Gene Names	Mme
Tag Info	quantitative
Protein Description	Sandwich
Description	This Mouse MME ELISA Kit was designed for the quantitative measurement of Mouse MME protein in serum, plasma, tissue homogenates. It is a Sandwich ELISA kit, its detection range is 0.156 ng/mL-10 ng/mL and the sensitivity is 0.039 ng/mL.
Target Details	This gene encodes a common acute lymphocytic leukemia antigen that is an

important cell surface marker in the diagnosis of human acute lymphocytic leukemia (ALL). This protein is present on leukemic cells of pre-B phenotype, which represent 85% of cases of ALL. This protein is not restricted to leukemic cells, however, and is found on a variety of normal tissues. It is a glycoprotein that is particularly abundant in kidney, where it is present on the brush border of

proximal tubules and on glomerular epithelium. The protein is a neutral

endopeptidase that cleaves peptides at the amino side of hydrophobic residues and inactivates several peptide hormones including glucagon, enkephalins, substance P, neurotensin, oxytocin, and bradykinin. This gene, which encodes a

CUSABIO® Your good partner in biology research



## **CUSABIO TECHNOLOGY LLC**





100-kD type II transmembrane glycoprotein, exists in a single copy of greater than 45 kb. The 5 untranslated region of this gene is alternatively spliced, resulting in four separate mRNA transcripts. The coding region is not affected by alternative splicing.

Product Precision	
Linearity	
Recovery	
Typical	
Msds	{"0":{"fileurl":"https://www.cusabio.com/uploadfile/msds/MSDS CSB-EL014653MO.pdf","filename":"MSDS"}}