



Human WNT1-inducible-signaling pathway protein 1(WISP1) ELISA kit

Product Code	CSB-EL026119HU
Abbreviation	WISP1
Protein Biological Process 1	Cell Adhesion
Target Name	WNT1 inducible signaling pathway protein 1
Uniprot No.	O95388
Alias	CCN4, WISP1c, WISP1i, WISP1tc, WNT1 induced secreted protein 1 Wnt-1 inducible signaling pathway protein 1 wnt-1 signaling pathway protein 1
Product Type	ELISA Kit
Immunogen Species	Homo sapiens (Human)
Protein Biological Process 3	Cell adhesion
Sample Types	serum, plasma, tissue homogenates
Detection Range	0.312 ng/mL-20 ng/mL
Sensitivity	0.078 ng/mL
Assay Time	1-5h
Sample Volume	50-100ul
Detection Wavelength	450 nm
Lead Time	3-5 working days after you place the order, and it takes another 3-5 days for delivery via DHL or FedEx.
Research Area	Epigenetics and Nuclear Signaling
Gene Names	WISP1
Tag Info	quantitative
Protein Description	Sandwich
Description	This Human WISP1 ELISA Kit was designed for the quantitative measurement of Human WISP1 protein in serum, plasma, tissue homogenates. It is a Sandwich ELISA kit, its detection range is 0.312 ng/mL-20 ng/mL and the sensitivity is 0.078 ng/mL.
Target Details	This gene encodes a member of the WNT1 inducible signaling pathway (WISP) protein subfamily, which belongs to the connective tissue growth factor (CTGF) family. WNT1 is a member of a family of cysteine-rich, glycosylated signaling proteins that mediate diverse developmental processes. The CTGF family



members are characterized by four conserved cysteine-rich domains: insulin-like growth factor-binding domain, von Willebrand factor type C module, thrombospondin domain and C-terminal cystine knot-like domain. This gene may be downstream in the WNT1 signaling pathway that is relevant to malignant transformation. It is expressed at a high level in fibroblast cells, and overexpressed in colon tumors. The encoded protein binds to decorin and biglycan, two members of a family of small leucine-rich proteoglycans present in the extracellular matrix of connective tissue, and possibly prevents the inhibitory activity of decorin and biglycan in tumor cell proliferation. It also attenuates p53-mediated apoptosis in response to DNA damage through activation of the Akt kinase. It is 83% identical to the mouse protein at the amino acid level. Alternative splicing of this gene generates 2 transcript variants.

Product Precision

Intra-assay Precision (Precision within an assay): CV%<8%

Three samples of known concentration were tested twenty times on one plate to assess.

Inter-assay Precision (Precision between assays): CV%<10%

Three samples of known concentration were tested in twenty assays to assess.

Linearity

To assess the linearity of the assay, samples were spiked with high concentrations of human WISP1 in various matrices and diluted with the Sample Diluent to produce samples with values within the dynamic range of the assay.

	Sample	Serum(n=4)
1:1	Average %	91
	Range %	86-95
1:2	Average %	102
	Range %	97-107
1:4	Average %	91
	Range %	85-97
1:8	Average %	97
	Range %	91-103


Recovery

The recovery of human WISP1 spiked to levels throughout the range of the assay in various matrices was evaluated. Samples were diluted prior to assay as directed in the Sample Preparation section.

Sample Type	Average % Recovery	Range
Serum (n=5)	95	89-98
EDTA plasma (n=4)	92	88-95

Typical

These standard curves are provided for demonstration only. A standard curve should be generated for each set of samples assayed.



ng/ml	OD1	OD2	Average	Corrected
20	2.058	2.162	2.110	1.937
10	1.654	1.620	1.637	1.464
5	1.241	1.298	1.270	1.097
2.5	0.834	0.864	0.849	0.676
1.25	0.512	0.532	0.522	0.349
0.625	0.344	0.361	0.353	0.180
0.312	0.242	0.262	0.252	0.079
0	0.172	0.174	0.173	

Msds

```
{
  "0": {
    "fileurl": "https://www.cusabio.com/uploadfile/msds/MSDS CSB-EL026119HU.pdf",
    "filename": "MSDS"
  }
}
```