




Recombinant Human Multidrug resistance-associated protein 1 (ABCC1), partial

Product Code	CSB-EP001056HU
Relevance	Mediates export of organic anions and drugs from the cytoplasm. Mediates ATP-dependent transport of glutathione and glutathione conjugates, leukotriene C4, estradiol-17-beta-o-glucuronide, methotrexate, antiviral drugs and other xenobiotics. Confers resistance to anticancer drugs. Hydrolyzes ATP with low efficiency.
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Uniprot No.	P33527
Product Type	Recombinant Protein
Immunogen Species	Homo sapiens (Human)
Purity	Greater than 90% as determined by SDS-PAGE.
Sequence	VRMSSEMETNIVAVERLKEYSETEKEAPWQIQETAPPSSWPQVGRVEFRNYC LRYREDLDFVLRHINVTINGGEKVGIVGRTGAGKSSLTLGLFRINESAEGEIIIDG INIAKIGLHDLRFKITIIPQDPVLFSGSLRMNLDPFQYSDEEVWTSLELAHLKDF VSALPDKLDHECAEGGENLSVGQRQLVCLARALLRKTILVLDEATAAVDLETD DLIQSTIRTQFEDCTVLTIAHRLNTIMDYTRVIVLDKGEIQEYGAPSDLLQQRGLF YSMA kDaGLV
Lead Time	3-7 business days
Research Area	Signal Transduction
Source	E.coli
Gene Names	ABCC1
Protein Names	ATP-binding cassette sub-family C member 1 Leukotriene C(4) transporter
Expression Region	1248-1531aa
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	N-terminal 6xHis-tagged
Mol. Weight	35.9 kDa
Protein Description	Partial
Image	



(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

Description

The production of this recombinant Human ABCC1 protein is just like all recombinant proteins. The process involved transfecting E.coli cells with DNA vector containing the template of recombinant DNA. The E.coli cells containing the template were then cultured so that they could transcribe and translate the ABCC1 protein. N-terminal 6xHis tag was used in the process. The purity is 0.9 determined by SDS-PAGE.

Multidrug resistance-associated protein 1 (MRP1/ABCC1) is a member of the ATP-binding cassette (ABC) transporter superfamily which contains 49 members in human that are divided into 7 subfamilies, named from ABCA to ABCG. (MRP1/ABCC1 is the first identified member of ABCC subfamily which belongs to ATP-binding cassette (ABC) transporter superfamily. It is ubiquitously expressed in almost all human tissues and transports a wide spectrum of substrates including drugs, heavy metal anions, toxicants, and conjugates of glutathione, glucuronide and sulfate. With the advance of sequence technology, many MRP1/ABCC1 polymorphisms have been identified. Accumulating evidences show that some polymorphisms are significantly associated with drug resistance and disease susceptibility. Recent advances showed the role and mechanism of MRP1/ABCC1 polymorphisms in drug resistance, toxicity, disease susceptibility and severity, prognosis prediction, and methods to select and predict functional polymorphisms.

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.