




Recombinant Human Aldo-keto reductase family 1 member C3 (AKR1C3)

Product Code	CSB-EP001544HU
Relevance	Catalyzes the conversion of aldehydes and ketones to alcohols. Catalyzes the reduction of prostaglandin (PG) D2, PGH2 and phenanthrenequinone (PQ) and the oxidation of 9-alpha,11-beta-PGF2 to PGD2. Functions as a bi-directional 3-alpha-, 17-beta- and 20-alpha HSD. Can interconvert active androgens, estrogens and progestins with their cognate inactive metabolites. Preferentially transforms androstenedione (4-dione) to testosterone.
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Uniprot No.	P42330
Alias	17-beta-hydroxysteroid dehydrogenase type 5
Product Type	Recombinant Protein
Immunogen Species	Homo sapiens (Human)
Purity	Greater than 90% as determined by SDS-PAGE.
Sequence	MDSKHQCVKLNLDGHFMPVLGFGTYAPPEVPRSKALEVTKLAIEAGFRHIDSAH LYNNEEQVGLAIRSKIADGSKREDIFYTSKLWSTFHRPELVRPALENSLKKQAQ LDYVDLYLIHSPMSLKPGEELSPTDENGKVIFDIVDLCTTWEAMEKCKDAGLAK SIGVSNFNRRQLEMILNKPGLKYKPVCNQVECHPYFNRSKLLDFCKSKDIVLVA YSALGSQRDKRWVDPNSPVLLEDPVLCALAKKHKRTPALIALRYQLQRGVVVL AKSYNEQRIRQNVQVFEFQLTAEDMKAIDGLDRNLHYFNSSDSFASHPNYPYSD EY
Lead Time	3-7 business days
Research Area	Signal Transduction
Source	E.coli
Gene Names	AKR1C3
Protein Names	Recommended name: Aldo-keto reductase family 1 member C3 EC= 1.-.-. Alternative name(s): 17-beta-hydroxysteroid dehydrogenase type 5 Short name= 17-beta-HSD 5 3-alpha-HSD type II, brain 3-alpha-hydroxysteroid dehydrogenase t
Expression Region	1-323aa
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	N-terminal GST-tagged
Mol. Weight	63.9kDa


Protein Description

Full Length

Image


(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.