




Recombinant Human Interferon gamma (IFNG)

Product Code	CSB-EP011050HU
Relevance	Produced by lymphocytes activated by specific antigens or mitogens. IFN-gamma, in addition to having antiviral activity, has important immunoregulatory functions. It is a potent activator of macrophages, it has antiproliferative effects on transformed cells and it can potentiate the antiviral and antitumor effects of the type I interferons.
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Uniprot No.	P01579
Alias	Immune interferon
Product Type	Recombinant Protein
Immunogen Species	Homo sapiens (Human)
Purity	Greater than 90% as determined by SDS-PAGE.
Sequence	QDPYVKEAENLKKYFNAGHSDVADNGTLFLGILKNWKEESDRKIMQSQIVSFY FKLFKNFKDDQSIQKSVETIKEDMNVKFFNSNKKKRDDFEKLTNYSVTDLNVQ RKAIHELIQVMAELSPAAGTGKRKRSQMLFRG
Lead Time	3-7 business days
Research Area	Immunology
Source	E.coli
Gene Names	IFNG
Expression Region	24-161aa
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	N-terminal 6xHis-tagged
Mol. Weight	20.2kDa
Protein Description	Full Length of Mature Protein
Image	



(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

Description

Cloning the gene encoding the Human IFNG protein (24-161aa) into a plasmid vector leads to the formation of recombinant plasmid. The resulting recombination plasmid is transformed into e.coli cells. e.coli cells containing the recombinant plasmid survive in the presence of a specific antibiotic and are selected to be cultured under conditions conducive to the expression of the gene of interest. A N-terminal 6xHis tag is linked to the protein. Following expression, the recombinant Human IFNG protein is isolated and purified from the cell lysate using affinity purification. Denaturing SDS-PAGE is utilized to resolve the resulting recombinant Human IFNG protein, demonstrating a purity exceeding 90%.

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.