



Recombinant Human Transmembrane protease serine 2 (TMPRSS2), partial

Product Code	CSB-EP023924HUe0
Relevance	Plasma membrane-anchored serine protease that participates in proteolytic cascades of relevance for the normal physiologic function of the prostate (PubMed:25122198). Androgen-induced TMPRSS2 activates several substrates that include pro-hepatocyte growth factor/HGF, the protease activated receptor-2/F2RL1 or matriptase/ST14 leading to extracellular matrix disruption and metastasis of prostate cancer cells (PubMed:15537383, PubMed:26018085, PubMed:25122198). In addition, activates trigeminal neurons and contribute to both spontaneous pain and mechanical allodynia
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Uniprot No.	O15393
Form	Liquid or Lyophilized powder
Alias	(Serine protease 10) (PRSS10)
Product Type	Recombinant Protein
Immunogen Species	Homo sapiens (Human)
Purity	Greater than 85% as determined by SDS-PAGE.
Sequence	WKFMGSKCSNSGIECDSSGTCINPSNWCDGVSHCPGGEDENRCVRLYGPNF ILQVYSSQRKSWHPVCQDDWNENYGRAACRDMGYKNNFYSSQGIVDDSGST SFMKLNTSAGNVDIYKKLYHSDACSSKAVVSLRCIACGVNLSSRQSRIVGGE SALPGAWPWQVSLHVQNVHVCSSITPEWIVTAAHCVEKPLNNPWHWTAF GILRQSFMFYAGYQVEKVISHPNYDSKTKNNDIALMKLQKPLTFNDLVKPVCL PNPGMMLQPEQLCWISGWGATEEKGKTSEVLNAAKVLLIETQRCNSRYVYDN LITPAMICAGFLQGNVDSCQGDSSGGLVTSKNNIWWLIGDTSWGSGCAKAYR PGVYGNVMVFTDWIYRQMRADG
Lead Time	Delivery time may differ from different purchasing way or location, please kindly consult your local distributors for specific delivery time.
Source	E.coli
Gene Names	TMPRSS2
Protein Names	Transmembrane protease serine 2
Expression Region	106-492 aa
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	N-terminal GST-tagged

**Protein Description**

Partial

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.