






## Recombinant Macaca fascicularis Interleukin-4 (IL4)

<b>Product Code</b>	CSB-EP301281MOV
Relevance	Participates in at least several B-cell activation processes as well as of other cell types. It is a costimulator of DNA-synthesis. It induces the expression of class II MHC molecules on resting B-cells. It enhances both secretion and cell surface expression of IgE and IgG1. It also regulates the expression of the low affinity Fc receptor for IgE (CD23) on both lymphocytes and monocytes. Positively regulates IL31RA expression in macrophages.
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Uniprot No.	P79339
Product Type	Recombinant Protein
Immunogen Species	Macaca fascicularis (Crab-eating macaque) (Cynomolgus monkey)
Purity	Greater than 85% as determined by SDS-PAGE.
Sequence	HKCDITLQEIIKTLNSLTEQKTLCTKLTITDILAASKNTTEKETFCRAATVLRQFYS HHEKDTRCLGATAQQFHRHKQLIRFLKRLDRNLWGLAGLNSCPVKEANQSTL ENFLERLKTIMREKYSKCSS
Lead Time	Delivery time may differ from different purchasing way or location, please kindly consult your local distributors for specific delivery time.
Research Area	Immunology
Source	E.coli
Gene Names	IL4
<b>Protein Names</b>	B-cell stimulatory factor 1 (BSF-1) (Lymphocyte stimulatory factor 1)
Expression Region	25-153aa
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	N-terminal 6xHis-tagged
Mol. Weight	18.9 kDa
<b>Protein Description</b>	Full Length of Mature Protein
Image	



## **CUSABIO TECHNOLOGY LLC**





(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

## Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL.We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.