




Recombinant Human RISC-loading complex subunit TARBP2 (TARBP2)

Product Code	CSB-EP623007HU
Relevance	Required for formation of the RNA induced silencing complex (RISC). Component of the RISC loading complex (RLC), also known as the micro-RNA (miRNA) loading complex (miRLC), which is composed of DICER1, AGO2 and TARBP2. Within the RLC/miRLC, DICER1 and TARBP2 are required to process precursor miRNAs (pre-miRNAs) to mature miRNAs and then load them onto AGO2. AGO2 bound to the mature miRNA constitutes the minimal RISC and may subsequently dissociate from DICER1 and TARBP2. May also play a role in the production of short interfering RNAs (siRNAs) from double-stranded RNA (dsRNA) by DICER1. Binds to the HIV-1 TAR RNA which is located in the long terminal repeat (LTR) of HIV-1, and stimulates translation of TAR-containing RNAs. This is achieved in part at least by binding to and inhibiting EIF2AK2/PKR, thereby reducing phosphorylation and inhibition of EIF2S1/eIF-2-alpha. May also promote translation of TAR-containing RNAs independently of EIF2AK2/PKR.
Abbreviation	TARBP2
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Uniprot No.	Q15633
Alias	TAR RNA-binding protein 2 Trans-activation-responsive RNA-binding protein
Product Type	Recombinant Protein
Immunogen Species	Homo sapiens (Human)
Purity	Greater than 90% as determined by SDS-PAGE.
Sequence	MSEEEQGS GTTTGCGLPSIEQMLAANPGKTPISLLQEYGTRIGKTPVYDLLKAE GQAHQPNFTFRVTVGDTSGTGQGPSKKA AKHKAAEVALKHLKGGSMLEPALE DSSSFSP LDSSLPEDIPVFTAAAAATPVPSVVLTRSPPMELQPPVSPQQSECN PVGALQELVVQKGWRLPEYTVTQESGPAHRKEFTMTCRVERFIEIGSGTSKKL AKRNAAAKMLLRVHTVPLDARDGNEVEPDDDDHFSIGVGSRLDGLRNRGPGCT WDSLRNSVGEKILSLRCSLGLSGALGPACCRVLSELSEEQAFHVS YLDIEELS LSGLCQCLVELSTQPATVCHGSATTREAAARGEAAARRALQYLKIMAGSK
Lead Time	Delivery time may differ from different purchasing way or location, please kindly consult your local distributors for specific delivery time.
Research Area	Microbiology
Source	E.coli
Gene Names	TARBP2



Protein Names	Recommended name: RISC-loading complex subunit TARBP2 Alternative name(s): TAR RNA-binding protein 2 Trans-activation-responsive RNA-binding protein
Expression Region	1-366aa
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	N-terminal GST-tagged
Mol. Weight	66.0kDa
Protein Description	Full Length

Image


(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.
-----------------------	---