




# Recombinant Mouse UMP-CMP kinase (Cmpk1)

|                            |  |
|----------------------------|--|
| <b>Product Code</b>        | CSB-EP875274MO   |
| <b>Relevance</b>           | Catalyzes the phosphorylation of pyrimidine nucleoside monophosphates at the expense of ATP. Plays an important role in de novo pyrimidine nucleotide biosynthesis. Has preference for UMP and CMP as phosphate acceptors. Also displays broad nucleoside diphosphate kinase activity. |
| <b>Storage</b>             | The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.        |
| <b>Uniprot No.</b>         | Q9DBP5   |
| <b>Alias</b>               | Deoxycytidylate kinaseU Short name: CK Short name: dCMP kinase<br>Nucleoside-diphosphate kinase Uridine monophosphate/cytidine monophosphate kinase Short name: UMP/CMP kinase Short name: UMP/CMP   |
| <b>Product Type</b>        | Recombinant Protein  |
| <b>Immunogen Species</b>   | Mus musculus (Mouse)   |
| <b>Purity</b>              | Greater than 90% as determined by SDS-PAGE.  |
| <b>Sequence</b>            | MKPLVVFVLGGPGAGKGTQCARIVEKYGYTHLSAGELLRDERKNPDSQYGELI<br>EKYIKEGKIVPVEITISLLKREMDQTMAANAQKNKFLIDGFPRNQDNLQGWNKT<br>MDGKADVSVLFFDCNNEICIERCLERKGSSGRSDDNRESLEKRIQTYLESTK<br>PIIDL YEEMGKVKKIDASKSVDEVFGEVVKIFDKEG   |
| <b>Lead Time</b>           | Delivery time may differ from different purchasing way or location, please kindly consult your local distributors for specific delivery time.  |
| <b>Research Area</b>       | Epigenetics and Nuclear Signaling  |
| <b>Source</b>              | E.coli   |
| <b>Gene Names</b>          | Cmpk1  |
| <b>Expression Region</b>   | 1-196aa  |
| <b>Notes</b>               | Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.  |
| <b>Tag Info</b>            | N-terminal 6xHis-tagged  |
| <b>Mol. Weight</b>         | 26.2kDa  |
| <b>Protein Description</b> | Full Length  |
| <b>Image</b>               |  |



(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

## Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.