





Recombinant Human mRNA-decapping enzyme 1A (DCP1A)

| Product Code | CSB-EP885684HU |
|---------------------|--|
| Relevance | Necessary for the degradation of mRNAs, both in normal mRNA turnover and in nonsense-mediated mRNA decay. Removes the 7-methyl guanine cap structure from mRNA molecules, yielding a 5'-phosphorylated mRNA fragment and 7m-GDP. Contributes to the transactivation of target genes after stimulation by TGFB1. |
| Abbreviation | DCP1A |
| Storage | The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C. |
| Uniprot No. | Q9NPI6 |
| Alias | Smad4-interacting transcriptional co-activator Transcription factor SMIF |
| Product Type | Recombinant Protein |
| Immunogen Species | Homo sapiens (Human) |
| Purity | Greater than 90% as determined by SDS-PAGE. |
| Sequence | MEALSRAGQEMSLAALKQHDPYITSIADLTGQVALYTFCPKANQWEKTDIEGT LFVYRRSASPYHGFTIVNRLNMHNLVEPVNKDLEFQLHEPFLLYRNASLSIYSI WFYDKNDCHRIAKLMADVVEEETRRSQQAARDKQSPSQANGCSDHRPIDILE MLSRAKDEYERNQMGDSNISSPGLQPSTQLSNLGSTETLEEMPSGSQDKSAP SGHKHLTVEELFGTSLPKEQPAVVGLDSEEMERLPGDASQKEPNSFLPFPFE QLGGAPQSETLGVPSAAHHSVQPEITTPVLITPASITQSNEKHAPTYTIPLSPVL SPTLPAEAPTAQVPPSLPRNSTMMQAVKTTPRQRSPLLNQPVPELSHASLIAN QSPFRAPLNVTNTAGTSLPSVDLLQKLRLTPQHDQIQTQPLGKGAMVASFSPA AGQLATPESFIEPPSKTAAARVAASASLSNMVLAPLQSMQQNQDPEVFVQPK VLSSAIQVAGAPLVTATTTAVSSVLLAPSVFQQTVTRSSDLERKASSPSPLTIGT PESQRKPSIILSKSQLQDTLIHLIKNDSSFLSTLHEVYLQVLTKNKDNHNL |
| Lead Time | Delivery time may differ from different purchasing way or location, please kindly consult your local distributors for specific delivery time. |
| Research Area | Epigenetics and Nuclear Signaling |
| Source | E.coli |
| Gene Names | DCP1A |
| Protein Names | Recommended name: mRNA-decapping enzyme 1A EC= 3 Alternative name(s): Smad4-interacting transcriptional co-activator Transcription factor SMIF |
| Expression Region | 1-582aa |
| | |



CUSABIO TECHNOLOGY LLC





Notes

Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.

Tag Info

N-terminal 6xHis-tagged


Mol. Weight

67.3kDa

Protein Description

Full Length

Image



(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL.We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.