




Recombinant Human 5'-AMP-activated protein kinase subunit gamma-2 (PRKAG2)

Product Code	CSB-EP887021HU
Relevance	AMP/ATP-binding subunit of AMP-activated protein kinase (AMPK), an energy sensor protein kinase that plays a key role in regulating cellular energy metabolism. In response to reduction of intracellular ATP levels, AMPK activates energy-producing pathways and inhibits energy-consuming processes: inhibits protein, carbohydrate and lipid biosynthesis, as well as cell growth and proliferation. AMPK acts via direct phosphorylation of metabolic enzymes, and by longer-term effects via phosphorylation of transcription regulators. Also acts as a regulator of cellular polarity by remodeling the actin cytoskeleton; probably by indirectly activating myosin. Gamma non-catalytic subunit mediates binding to AMP, ADP and ATP, leading to activate or inhibit AMPK: AMP-binding results in allosteric activation of alpha catalytic subunit (PRKAA1 or PRKAA2) both by inducing phosphorylation and preventing dephosphorylation of catalytic subunits. ADP also stimulates phosphorylation, without stimulating already phosphorylated catalytic subunit. ATP promotes dephosphorylation of catalytic subunit, rendering the AMPK enzyme inactive.
Abbreviation	PRKAG2
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Uniprot No.	Q9UGJ0
Alias	H91620p
Product Type	Recombinant Protein
Immunogen Species	Homo sapiens (Human)
Purity	Greater than 90% as determined by SDS-PAGE.
Sequence	MLEKLEFEDEAVEDSESGVYMRFMRSCHKCYDIVPTSSKLVVFDTTLQVKKAFF ALVANGVRAAPLWESKKQSFVGMTITDFINILHRYKSPMVQIYELEEHKIET WRELYLQETFKPLVNISPDASLFDVYSLIKNIHRLPVIDPISGNALYILTHKRIL KFLQLFMSDMPKPAFMKQNLDELIGITYHNIAFIHPDTPIIKALNIFVERRISALP VVDESGKVVDIYSKFDVINLAAEKTNNLDITVTQALQHRSQYFEGVVKCNKLEI LETIVDRIVRAEVHRLVVVNEADSIVGIISLSDILQALILTPAGAKQKETETE
Lead Time	Delivery time may differ from different purchasing way or location, please kindly consult your local distributors for specific delivery time.
Research Area	Cardiovascular
Source	E.coli
Gene Names	PRKAG2



Protein Names	Recommended name: 5'-AMP-activated protein kinase subunit gamma-2 Short name= AMPK gamma2 Short name= AMPK subunit gamma-2Alternative name(s): H91620p
Expression Region	1-328aa
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	N-terminal 6xHis-tagged
Mol. Weight	41.5kDa
Protein Description	Full Length of Isoform B

Image


(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.
-----------------------	---