



Recombinant Human Nuclear pore complex protein Nup98-Nup96 (NUP98), partial

Product Code	CSB-MP016209HU
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Uniprot No.	P52948
Form	Liquid or Lyophilized powder
Storage Buffer	If the delivery form is liquid, the default storage buffer is Tris/PBS-based buffer, 5%-50% glycerol. If the delivery form is lyophilized powder, the buffer before lyophilization is Tris/PBS-based buffer, 6% Trehalose.
Product Type	Recombinant Protein
Immunogen Species	Homo sapiens (Human)
Purity	Greater than 85% as determined by SDS-PAGE.
Sequence	MFNKSFGTPFGGGTGGFGTTSTFGQNTGFGTSSGGAFGTSAFGSSNNTGGL FGNSQTKPGGLFGTSSFSQPATSTSTGFGGTSTGTANTLFGTASTGTSLFSS QNNFAQNKPTGFGNFGTSTSSGGLFGTTNTTSNPFGSTSGSLFGPSSFTAA PTGTTIKFNPPTGDTMVKAGVSTNISTKHQCITAMKEYESKSLEELRLEDYQA NRKGPQNQVGAGTTTGLFGSSPATSSATGLFSSSTNSGFAYGQNKTAFGTS TTGFGTNPGGFLGQQNQQTTSLSKPFQATTTQNTGFSFGNTSTIGQPSTN TMGLFGVTQASQPGGLFGTATNTSTGTAFGTGTGLFGQTNTGFGAVGSTLFG NNKLTTFGSSTTSAPSFGTSSGGLFGNKPTLTLGTNTNTSNFGFGTNTSGNSIF GSKPAPGTLGTGLGAGFGTALGAGQASLFGNNQPKIGGPLGTGAFGAPGFNT TTATLFGGAPQAPVALTDPNASAAQQAVALQQHINSLTYSFPGDSPLFRNPMSD PKKKEERLKPTNPAAQKALTTPTHYKLTTPRPA TRVRPKALQTTGTAKSHLFDG LDDDEPSLANGAFMPKKSIIKKLVKLNLSNLFSPVNRDSENLASPSEYPENG ERFSFLSKPVDENHQDGEDSLVSHFYTNPIAKPIQTPESAGNKHNSNSV DDTIVALNMRAALRNGLEGSSEETSFHDESLQDDREEIENNSYHMHAPAGIILTK VGYYTIPSMDDLAKITNEKGECIVSDFTIGRKGYGSIYFEGDVNLTNLNLDDIVHI RRKEVVVYLDDNQKPPVGEGLNRKAEVTLDG VWP TDKTSRCLIKSPDRLADIN YEGRLEAVSRKQGAQFKEYRPETGSWVFKVSHF
Lead Time	3-7 business days
Research Area	Epigenetics and Nuclear Signaling
Source	Mammalian cell
Gene Names	NUP98
Expression Region	1-880aa
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.




Tag Info C-terminal hFc-tagged

Mol. Weight 120.6 kDa

Protein Description Partial

Image



(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.