





## Recombinant Escherichia coli tRNA threonylcarbamoyladenosine biosynthesis protein TsaE (tsaE)

Product Code	CSB-MP360227ENV
Relevance	Required for the formation of a threonylcarbamoyl group on adenosine at position 37 (t6A37) in tRNAs that read codons beginning with adenine. Is probably involved in the transfer of the threonylcarbamoyl moiety of threonylcarbamoyl-AMP (TC-AMP) to the N6 group of A37, together with TsaD and TsaB. TsaE seems to play an indirect role in the t6A biosynthesis pathway, possibly in regulating the core enzymatic function of TsaD. Displays ATPase activity in vitro.
Storage	The shelf life is related to many factors, storage state, buffer ingredients, storage temperature and the stability of the protein itself. Generally, the shelf life of liquid form is 6 months at -20°C/-80°C. The shelf life of lyophilized form is 12 months at -20°C/-80°C.
Uniprot No.	P0AF67
Alias	t(6)A37 threonylcarbamoyladenosine biosynthesis protein TsaE
Product Type	Recombinant Protein
Immunogen Species	Escherichia coli (strain K12)
Purity	Greater than 90% as determined by SDS-PAGE.
Sequence	MMNRVIPLPDEQATLDLGERVAKACDGATVIYLYGDLGAGKTTFSRGFLQALG HQGNVKSPTYTLVEPYTLDNLMVYHFDLYRLADPEELEFMGIRDYFANDAICLV EWPQQGTGVLPDPDVEIHIDYQAQGREARVSAVSSAGELLLARLAG
Lead Time	Delivery time may differ from different purchasing way or location, please kindly consult your local distributors for specific delivery time.
Research Area	Others
Source	Mammalian cell
Gene Names	tsaE
Protein Names	Recommended name: UPF0079 ATP-binding protein yjeE
Expression Region	1-153aa
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.
Tag Info	C-terminal Flag-tagged
Mol. Weight	27.5kDa
<b>Protein Description</b>	Full Length
Image	




## **CUSABIO TECHNOLOGY LLC**

🕜 Tel: +1-301-363-4651 🛛 Email: cusabio@cusabio.com 🕒 Website: www.cusabio.com 🌘







(Tris-Glycine gel) Discontinuous SDS-PAGE (reduced) with 5% enrichment gel and 15% separation gel.

## Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL.We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.