






Phospho-CEBPB (T235) Antibody

Product Code CSB-PA000487 Storage Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. Uniprot No. P17676 Immunogen Synthesized peptide derived from Human C/EBP β around the phosphorylatisite of T235. Raised In Rabbit Species Reactivity Human,Mouse,Rat Tested Applications WB, IHC, IF, ELISA; WB:1:500-1:2000, IHC:1:100-1:300, IF:1:200-1:1000, ELISA:1:10000 Form Liquid Conjugate Non-conjugated Storage Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Purification Method The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Isotype IgG
Uniprot No.P17676ImmunogenSynthesized peptide derived from Human C/EBP β around the phosphorylating site of T235.Raised InRabbitSpecies ReactivityHuman,Mouse,RatTested ApplicationsWB, IHC, IF, ELISA; WB:1:500-1:2000, IHC:1:100-1:300, IF:1:200-1:1000, ELISA:1:10000FormLiquidConjugateNon-conjugatedStorage BufferLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.Purification MethodThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
ImmunogenSynthesized peptide derived from Human C/EBP β around the phosphorylating site of T235.Raised InRabbitSpecies ReactivityHuman,Mouse,RatTested ApplicationsWB, IHC, IF, ELISA; WB:1:500-1:2000, IHC:1:100-1:300, IF:1:200-1:1000, ELISA:1:10000FormLiquidConjugateNon-conjugatedStorage BufferLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.Purification MethodThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Raised In Rabbit Species Reactivity Human,Mouse,Rat Tested Applications WB, IHC, IF, ELISA; WB:1:500-1:2000, IHC:1:100-1:300, IF:1:200-1:1000, ELISA:1:10000 Form Liquid Conjugate Non-conjugated Storage Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Purification Method The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Species ReactivityHuman,Mouse,RatTested ApplicationsWB, IHC, IF, ELISA; WB:1:500-1:2000, IHC:1:100-1:300, IF:1:200-1:1000, ELISA:1:10000FormLiquidConjugateNon-conjugatedStorage BufferLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.Purification MethodThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Tested Applications WB, IHC, IF, ELISA; WB:1:500-1:2000, IHC:1:100-1:300, IF:1:200-1:1000, ELISA:1:10000 Form Liquid Non-conjugated Storage Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Purification Method The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Form Liquid Conjugate Non-conjugated Storage Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Purification Method The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Conjugate Non-conjugated Storage Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Purification Method The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Storage Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Purification Method The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Purification Method The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
chromatography using epitope-specific immunogen.
Isotype IaG
isotype igo
Alias CEBPB; LAP; TCF5; PP9092; CCAAT/enhancer-binding protein beta; C/EBF beta; Liver activator protein; Nuclear factor NF-IL6; Transcription factor 5; TC
Immunogen Species Homo sapiens (Human)
Target Names CEBPB
Image Western Blot analysis of HanG2 cells using



Western Blot analysis of HepG2 cells using Phospho-C/EBP β (T235) Polyclonal Antibody