




ERCC2 Antibody

Product Code	CSB-PA007770ESR2HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P18074
Immunogen	Recombinant Human TFIIH basal transcription factor complex helicase XPD subunit protein (1-300AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, WB, IHC; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Purification Method	Antigen Affinity Purified
Isotype	IgG
Clonality	Polyclonal
Alias	General transcription and DNA repair factor IIH helicase subunit XPD (TFIIH subunit XPD) (EC 3.6.4.12) (Basic transcription factor 2 80 kDa subunit) (BTF2 p80) (CXPB) (DNA excision repair protein ERCC-2) (DNA repair protein complementing XP-D cells) (TFIIH basal transcription factor complex 80 kDa subunit) (TFIIH 80 kDa subunit) (TFIIH p80) (Xeroderma pigmentosum group D-complementing protein), ERCC2, XPD XPDC
Immunogen Species	Homo sapiens (Human)
Research Area	Epigenetics and Nuclear Signaling
Target Names	ERCC2

Image



Western blot

All lanes: TFIIH basal transcription factor complex helicase XPD subunit antibody at 2μg/ml

Lane 1: 293T whole cell lysate


Lane 2: K562 whole cell lysate

Secondary


Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 87, 47 kDa

Observed band size: 87 kDa



Immunohistochemistry of paraffin-embedded human skin tissue using CSB-PA007770ESR2HU at dilution of 1:100



Immunohistochemistry of paraffin-embedded human lung cancer using CSB-PA007770ESR2HU at dilution of 1:100