






## FOLR1 Antibody

<b>Product Code</b>	CSB-PA008784LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P15328
Immunogen	Recombinant Human Folate receptor alpha protein (25-234AA)
Raised In	Rabbit
Species Reactivity	Human
<b>Tested Applications</b>	ELISA, IHC, IF; Recommended dilution: IHC:1:500-1:1000, IF:1:200-1:500
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
<b>Purification Method</b>	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	Folate receptor alpha (FR-alpha) (Adult folate-binding protein) (FBP) (Folate receptor 1) (Folate receptor, adult) (KB cells FBP) (Ovarian tumor-associated antigen MOv18), FOLR1, FOLR
Immunogen Species	Homo sapiens (Human)
Research Area	Tags & Cell Markers
Target Names	FOLR1
Image	IHC image of CSR DA009794LA01HI Lidiluted at



IHC image of CSB-PA008784LA01HU diluted at 1:800 and staining in paraffin-embedded human placenta tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.




## **CUSABIO TECHNOLOGY LLC**









Immunofluorescence staining of HepG2 cells with CSB-PA008784LA01HU at 1:266, counterstained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).