


Image





LAG3 Antibody

Product Code	CSB-PA012719LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P18627
Immunogen	Recombinant Human Lymphocyte activation gene 3 protein (29-450AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, WB, IF; Recommended dilution: WB:1:500-1:5000, IF:1:50-1:200
Form	Liquid
Conjugate	Non-conjugated
Storago Buffor	Preservative: 0.03% Proclin 300
Storage Buffer	Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Purification Method	
	Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Purification Method	Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified
Purification Method Isotype	Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG
Purification Method Isotype Clonality	Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG Polyclonal
Purification Method Isotype Clonality Product Type	Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG Polyclonal Polyclonal Antibody
Purification Method Isotype Clonality Product Type Immunogen Species	Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified IgG Polyclonal Polyclonal Antibody Homo sapiens (Human)



Western Blot

Positive WB detected in: HepG2 whole cell

All lanes: LAG3 antibody at 4.9µg/ml

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 58, 40 kDa Observed band size: 58 kDa




CUSABIO TECHNOLOGY LLC











Immunofluorescence staining of HepG2 cells with CSB-PA012719LA01HU at 1:66, counterstained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).