




# NTN1 Antibody

|                            |  |
|----------------------------|--|
| <b>Product Code</b>        | CSB-PA016127LA01HU   |
| <b>Storage</b>             | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.                    |
| <b>Uniprot No.</b>         | O95631   |
| <b>Immunogen</b>           | Recombinant Human Netrin-1 protein (186-478AA)                                   |
| <b>Raised In</b>           | Rabbit   |
| <b>Species Reactivity</b>  | Human  |
| <b>Tested Applications</b> | ELISA, WB, IHC; Recommended dilution: WB:1:500-1:5000, IHC:1:200-1:500           |
| <b>Form</b>                | Liquid   |
| <b>Conjugate</b>           | Non-conjugated   |
| <b>Storage Buffer</b>      | Preservative: 0.03% Proclin 300<br>Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 |
| <b>Purification Method</b> | >95%, Protein G purified   |
| <b>Isotype</b>             | IgG  |
| <b>Clonality</b>           | Polyclonal   |
| <b>Alias</b>               | Netrin-1 (Epididymis tissue protein Li 131P), NTN1, NTN1L                        |
| <b>Immunogen Species</b>   | Homo sapiens (Human)   |
| <b>Research Area</b>       | Neuroscience   |
| <b>Target Names</b>        | NTN1   |

## Image




### Western Blot

Positive WB detected in: 293T whole cell lysate, Hela whole cell lysate, HepG2 whole cell lysate  
All lanes: NTN1 antibody at 6.4µg/ml

### Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution  
Predicted band size: 68 kDa  
Observed band size: 68 kDa



IHC image of CSB-PA016127LA01HU diluted at 1:400 and staining in paraffin-embedded human liver tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a



biotinylated secondary antibody and visualized using an HRP conjugated SP system.

---