






## **NUCB2** Antibody

Product Code	CSB-PA016146HA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P80303
Immunogen	Recombinant Human Nucleobindin-2 protein (25-106AA)
Raised In	Rabbit
Species Reactivity	Human, Mouse, Rat
<b>Tested Applications</b>	ELISA, WB, IHC; Recommended dilution: WB:1:500-1:5000, IHC:1:20-1:200
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
<b>Purification Method</b>	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	Nucleobindin-2 (DNA-binding protein NEFA) (Gastric cancer antigen Zg4) (Prepronesfatin) [Cleaved into: Nesfatin-1], NUCB2, NEFA
Immunogen Species	Homo sapiens (Human)
Research Area	Signal Transduction
Target Names	NUCB2
les a se	





Western Blot

Positive WB detected in: Rat brain tissue, Rat heart tissue, Mouse heart tissue, Mouse liver

tissue, Mouse brain tissue

All lanes: NUCB2 antibody at 2.5µg/ml

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 51 kDa Observed band size: 51 kDa















IHC image of CSB-PA016146HA01HU diluted at 1:200 and staining in paraffin-embedded human small intestine tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC image of CSB-PA016146HA01HU diluted at 1:200 and staining in paraffin-embedded human lung cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.