











PRG2 Antibody

Product Code	CSB-PA018670LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P13727
Immunogen	Recombinant Human Bone marrow proteoglycan protein (151-195AA)
Raised In	Rabbit
Species Reactivity	Human, Mouse
Tested Applications	ELISA, WB, IHC; Recommended dilution: WB:1:1000-1:5000, IHC:1:50-1:000
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.05% NaN3 Constituents: PBS containing 40% glycerol pH7.4
Purification Method	Antigen affinity purification
Isotype	IgG
Clonality	Polyclonal
Product Type	Polyclonal
Immunogen Species	Homo sapiens (Human)
Research Area	Tags & Cell Markers
Target Names	PRG2
Image	ILIC image of CSR DA049670LA04LILI diluted at



IHC image of CSB-PA018670LA01HU diluted at 1:66 and staining in paraffin-embedded human liver cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.05% DAB. Secondary antibody only control: uses 1% BSA instead of primary antibody




IHC image of CSB-PA018670LA01HU diluted at 1:66 and staining in paraffin-embedded human breast cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat







anti-rabbit polymer IgG labeled by HRP and visualized using 0.05% DAB. Secondary antibody only control: uses 1% BSA instead of primary antibody



Western Blot

Positive WB detected in: Mouse spleen tissue

lysate, Rat spleen tissue lysate All lanes: PRG2 antibody at 1:1000

Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 25, 28 kDa Observed band size: 28 kDa