




PRPF40A Antibody

Product Code	CSB-PA018769LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	O75400
Immunogen	Recombinant Human Pre-mRNA-processing factor 40 homolog A protein (120-266AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, WB, IHC; Recommended dilution: WB:1:500-1:5000, IHC:1:200-1:500
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Purification Method	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	Pre-mRNA-processing factor 40 homolog A, Fas ligand-associated factor 1, Formin-binding protein 11, Formin-binding protein 3, Huntingtin yeast partner A, Huntingtin-interacting protein 10, HIP-10, Huntingtin-interacting protein A, Renal carcinoma antigen NY-REN-6, PRPF40A, FBP11, FLAF1, FNBP3, HIP10, HYPA, HSPC225
Immunogen Species	Homo sapiens (Human)
Research Area	Epigenetics and Nuclear Signaling
Target Names	PRPF40A

Image



Western Blot

Positive WB detected in: U251 whole cell lysate, HeLa whole cell lysate, 293T whole cell lysate


All lanes: PRPF40A antibody at 2.7µg/ml

Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 109, 106, 107, 11 kDa

Observed band size: 109 kDa



IHC image of CSB-PA018769LA01HU diluted at 1:200 and staining in paraffin-embedded human testis tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.