
**Image** 





## **TAAR5** Antibody

<b>Product Code</b>	CSB-PA023057LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	O14804
Immunogen	Recombinant Human Trace amine-associated receptor 5 protein (1-34AA)
Raised In	Rabbit
Species Reactivity	Human, Rat, Mouse
Tested Applications	ELISA, WB, IHC, IF; Recommended dilution: WB:1:500-1:5000, IHC:1:500-1:1000, IF:1:200-1:500
Form	Liquid
Conjugate	Non-conjugated
	December 1997 December 200
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Storage Buffer Purification Method	
	Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Purification Method	Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 >95%, Protein G purified
Purification Method Isotype	Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 >95%, Protein G purified  IgG
Purification Method Isotype Clonality	Constituents: 50% Glycerol, 0.01M PBS, pH 7.4  >95%, Protein G purified  IgG  Polyclonal  Trace amine-associated receptor 5 (TaR-5) (Trace amine receptor 5) (hTaar5)
Purification Method Isotype Clonality Alias	Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 >95%, Protein G purified  IgG  Polyclonal  Trace amine-associated receptor 5 (TaR-5) (Trace amine receptor 5) (hTaar5) (Putative neurotransmitter receptor), TAAR5, PNR
Purification Method Isotype Clonality Alias Immunogen Species	Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 >95%, Protein G purified  IgG  Polyclonal  Trace amine-associated receptor 5 (TaR-5) (Trace amine receptor 5) (hTaar5) (Putative neurotransmitter receptor), TAAR5, PNR  Homo sapiens (Human)



Western Blot

Positive WB detected in: Rat brain tissue, Mouse

brain tissue

All lanes: TAAR5 antibody at 3μg/ml


Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 39 kDa Observed band size: 39 kDa















IHC image of CSB-PA023057LA01HU diluted at 1:600 and staining in paraffin-embedded human brain tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunofluorescence staining of Hela cells with CSB-PA023057LA01HU at 1:200, counterstained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).