





# TCF12 Antibody

<b>Product Code</b>	CSB-PA023293ESR1HU
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	Q99081
<b>Immunogen</b>	Recombinant Human Transcription factor 12 protein (443-682AA)
<b>Raised In</b>	Rabbit
<b>Species Reactivity</b>	Human
<b>Tested Applications</b>	ELISA, IHC, IP; Recommended dilution: IHC:1:20-1:200, IP:1:200-1:2000
<b>Form</b>	Liquid
<b>Conjugate</b>	Non-conjugated
<b>Storage Buffer</b>	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
<b>Purification Method</b>	Antigen Affinity Purified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Alias</b>	Transcription factor 12 (TCF-12) (Class B basic helix-loop-helix protein 20) (bHLHb20) (DNA-binding protein HTF4) (E-box-binding protein) (Transcription factor HTF-4), TCF12, BHLHB20 HEB HTF4
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Research Area</b>	Epigenetics and Nuclear Signaling
<b>Target Names</b>	TCF12

## Image



Immunohistochemistry of paraffin-embedded human thymus tissue using CSB-PA023293ESR1HU at dilution of 1:100



Immunoprecipitating TCF12 in Jurkat whole cell lysate

Lane 1: Rabbit control IgG instead of (1 $\mu$ g) instead of CSB-PA023293ESR1HU in Jurkat whole cell lysate. For western blotting, a HRP-conjugated anti-rabbit IgG, specific to the non-reduced form of IgG was used as the Secondary antibody (1/50000)

Lane 2: CSB-PA023293ESR1HU (4 $\mu$ g) + Jurkat whole cell lysate (500 $\mu$ g)

Lane 3: Jurkat whole cell lysate (20 $\mu$ g)