




TGFBR1 Antibody

Product Code	CSB-PA023451LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	P36897
Immunogen	Synthetic peptide of human TGFBR1
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, WB, IHC, IF; Recommended dilution: WB:1:500-1:2000, IHC:1:100-1:300, IF:1:10-1:100
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	pH7.4 PBS, 0.05% NaN ₃ , 40% Glycerol
Purification Method	Antigen affinity purification
Isotype	IgG
Clonality	Polyclonal
Product Type	Polyclonal Antibody
Immunogen Species	Homo sapiens (Human)
Research Area	Cardiovascular
Target Names	TGFBR1

Image




Western Blot


Positive WB detected in: HepG2 whole cell lysate, A549 whole cell lysate, HeLa whole cell lysate, SH-SY5Y, PC-3 whole cell lysate
All lanes: TGFBR1 antibody at 1:1000

Secondary


Goat polyclonal to rabbit IgG at 1/50000 dilution
Predicted band size: 56, 57, 48kDa
Observed band size: 58, 40kDa



IHC image of CSB-PA023451LA01HU diluted at 1:1200 and staining in paraffin-embedded human small intestine tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunofluorescence staining of HepG2 cell with CSB-PA023451LA01HU at 1:40, counter-stained with DAPI. The cells were fixed in 4% formaldehyde and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescence staining of HepG2 cell with 5% goat serum, counter-stained with DAPI. The cells were fixed in 4% formaldehyde and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).