






## **UBE2V2** Antibody

Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.  Uniprot No. Q15819  Recombinant Human Ubiquitin-conjugating enzyme E2 variant 2 protein (2-145AA)  Raised In Rabbit  Human  Ested Applications  ELISA, IHC; Recommended dilution: IHC:1:20-1:200  Form Liquid  Non-conjugated  Non-conjugated  Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method >95%, Protein G purified  Sotype IgG  Clonality Polyclonal  Ubiquitin-conjugating enzyme E2 variant 2 (DDVit 1) (Enterocyte differentiation-associated factor 1) (EDPF-1) (MMS2 homolog) (Vitamin D3-inducible protein), UBE2V2, MMS2 UEV2  mmunogen Species Homo sapiens (Human)  Research Area Epigenetics and Nuclear Signaling  Garget Names UBE2V2		
Iniprot No. Q15819  Recombinant Human Ubiquitin-conjugating enzyme E2 variant 2 protein (2-145AA)  Raised In Rabbit  Recides Reactivity Human  Fested Applications ELISA, IHC; Recommended dilution: IHC:1:20-1:200  Form Liquid  Ronjugate Non-conjugated  Rorage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method >95%, Protein G purified  Sotype IgG  Clonality Polyclonal  Ubiquitin-conjugating enzyme E2 variant 2 (DDVit 1) (Enterocyte differentiation-associated factor 1) (EDAF-1) (Enterocyte differentiation-promoting factor 1) (EDF-1) (MMS2 homolog) (Vitamin D3-inducible protein), UBE2V2, MMS2 UEV2  Immunogen Species Homo sapiens (Human)  Research Area Epigenetics and Nuclear Signaling  Figure 1.	Product Code	CSB-PA025484HA01HU
Recombinant Human Ubiquitin-conjugating enzyme E2 variant 2 protein (2-145AA)  Raised In Rabbit  Recies Reactivity Human  Ested Applications ELISA, IHC; Recommended dilution: IHC:1:20-1:200  Form Liquid  Conjugate Non-conjugated  Storage Buffer Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method >95%, Protein G purified  Sotype IgG  Clonality Polyclonal  Ubiquitin-conjugating enzyme E2 variant 2 (DDVit 1) (Enterocyte differentiation-associated factor 1) (EDAF-1) (Enterocyte differentiation-promoting factor 1) (EDPF-1) (MMS2 homolog) (Vitamin D3-inducible protein), UBE2V2, MMS2  UEV2  Immunogen Species Homo sapiens (Human)  Research Area Epigenetics and Nuclear Signaling  Target Names UBE2V2	Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
(2-145AA)  Rabbit  Rabbit  Human  Elseted Applications  ELISA, IHC; Recommended dilution: IHC:1:20-1:200  Form  Liquid  Conjugate  Non-conjugated  Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method  Sotype  IgG  Clonality  Polyclonal  Ubiquitin-conjugating enzyme E2 variant 2 (DDVit 1) (Enterocyte differentiation-associated factor 1) (EDAF-1) (Enterocyte differentiation-promoting factor 1) (EDPF-1) (MMS2 homolog) (Vitamin D3-inducible protein), UBE2V2, MMS2  UEV2  Immunogen Species  Homo sapiens (Human)  Research Area  Epigenetics and Nuclear Signaling  Target Names  UBE2V2	Uniprot No.	Q15819
Figure Reactivity Figure Reservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 Purification Method >95%, Protein G purified  Figure Reservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method >95%, Protein G purified  Figure Reservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method >95%, Protein G purified  Figure Reservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method >95%, Protein G purified  Figure Reservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method >95%, Protein G purified  Figure Reservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method >95%, Protein G purified  Figure Reservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method >95%, Protein G purified  Figure Reservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method >95%, Protein G purified  Figure Reservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method >95%, Protein G purified  Figure Reservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method >95%, Protein G purified  Figure Reservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method >95%, Protein G purified  Figure Reservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method >95%, Protein G purified  Figure Reservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method >95%, Protein G purified  Figure Reservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method >95%, Protein G purified  Figure Reservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method >95%, Protein G purified  Figure Reservative: 0.03% Proclin 300 Constituents: 50% Glycerol	Immunogen	
ELISA, IHC; Recommended dilution: IHC:1:20-1:200  Form Liquid  Non-conjugated  Non-conjugated  Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method >95%, Protein G purified  Sotype IgG  Clonality Polyclonal  Ubiquitin-conjugating enzyme E2 variant 2 (DDVit 1) (Enterocyte differentiation-associated factor 1) (EDAF-1) (Enterocyte differentiation-promoting factor 1) (EDPF-1) (MMS2 homolog) (Vitamin D3-inducible protein), UBE2V2, MMS2 UEV2  Immunogen Species Homo sapiens (Human)  Research Area Epigenetics and Nuclear Signaling  Target Names UBE2V2	Raised In	Rabbit
Liquid  Non-conjugate Non-conjugated  Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method  >95%, Protein G purified  sotype  IgG  Clonality Polyclonal  Ubiquitin-conjugating enzyme E2 variant 2 (DDVit 1) (Enterocyte differentiation-associated factor 1) (EDAF-1) (Enterocyte differentiation-promoting factor 1) (EDPF-1) (MMS2 homolog) (Vitamin D3-inducible protein), UBE2V2, MMS2 UEV2  mmunogen Species  Homo sapiens (Human)  Research Area Epigenetics and Nuclear Signaling  Target Names  UBE2V2	Species Reactivity	Human
Non-conjugated  Non-conjugated  Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method >95%, Protein G purified  Sotype IgG  Clonality Polyclonal  Ubiquitin-conjugating enzyme E2 variant 2 (DDVit 1) (Enterocyte differentiation-associated factor 1) (EDAF-1) (Enterocyte differentiation-promoting factor 1) (EDPF-1) (MMS2 homolog) (Vitamin D3-inducible protein), UBE2V2, MMS2 UEV2  Immunogen Species Homo sapiens (Human)  Research Area Epigenetics and Nuclear Signaling  Target Names UBE2V2	<b>Tested Applications</b>	ELISA, IHC; Recommended dilution: IHC:1:20-1:200
Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method >95%, Protein G purified  Sotype IgG  Clonality Polyclonal  Ubiquitin-conjugating enzyme E2 variant 2 (DDVit 1) (Enterocyte differentiation-associated factor 1) (EDAF-1) (Enterocyte differentiation-promoting factor 1) (EDPF-1) (MMS2 homolog) (Vitamin D3-inducible protein), UBE2V2, MMS2 UEV2  Immunogen Species Homo sapiens (Human)  Research Area Epigenetics and Nuclear Signaling  Garget Names UBE2V2	Form	Liquid
Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  Purification Method >95%, Protein G purified  sotype IgG  Clonality Polyclonal  Ubiquitin-conjugating enzyme E2 variant 2 (DDVit 1) (Enterocyte differentiation-associated factor 1) (EDAF-1) (Enterocyte differentiation-promoting factor 1) (EDPF-1) (MMS2 homolog) (Vitamin D3-inducible protein), UBE2V2, MMS2 UEV2  mmunogen Species Homo sapiens (Human)  Research Area Epigenetics and Nuclear Signaling  Target Names UBE2V2	Conjugate	Non-conjugated
IgG  Clonality Polyclonal  Ubiquitin-conjugating enzyme E2 variant 2 (DDVit 1) (Enterocyte differentiation-associated factor 1) (EDAF-1) (Enterocyte differentiation-promoting factor 1) (EDPF-1) (MMS2 homolog) (Vitamin D3-inducible protein), UBE2V2, MMS2 UEV2  mmunogen Species Homo sapiens (Human)  Research Area Epigenetics and Nuclear Signaling  Target Names UBE2V2	Storage Buffer	
Clonality Polyclonal Ubiquitin-conjugating enzyme E2 variant 2 (DDVit 1) (Enterocyte differentiation-associated factor 1) (EDAF-1) (Enterocyte differentiation-promoting factor 1) (EDPF-1) (MMS2 homolog) (Vitamin D3-inducible protein), UBE2V2, MMS2 UEV2  mmunogen Species Homo sapiens (Human)  Research Area Epigenetics and Nuclear Signaling Target Names UBE2V2	Purification Method	>95%, Protein G purified
Ubiquitin-conjugating enzyme E2 variant 2 (DDVit 1) (Enterocyte differentiation-associated factor 1) (EDAF-1) (Enterocyte differentiation-promoting factor 1) (EDPF-1) (MMS2 homolog) (Vitamin D3-inducible protein), UBE2V2, MMS2 UEV2  mmunogen Species Homo sapiens (Human)  Research Area Epigenetics and Nuclear Signaling  Target Names UBE2V2	Isotype	IgG
associated factor 1) (EDAF-1) (Enterocyte differentiation-promoting factor 1) (EDPF-1) (MMS2 homolog) (Vitamin D3-inducible protein), UBE2V2, MMS2 UEV2  mmunogen Species Homo sapiens (Human)  Research Area Epigenetics and Nuclear Signaling  Target Names UBE2V2  mage	Clonality	Polyclonal
Research Area Epigenetics and Nuclear Signaling Target Names UBE2V2 mage	Alias	associated factor 1) (EDAF-1) (Enterocyte differentiation-promoting factor 1) (EDPF-1) (MMS2 homolog) (Vitamin D3-inducible protein), UBE2V2, MMS2
Target Names UBE2V2	Immunogen Species	Homo sapiens (Human)
mage	Research Area	Epigenetics and Nuclear Signaling
mage	Target Names	UBE2V2
	Image	Immunohistochemistry of paraffin-embedded



Immunohistochemistry of paraffin-embedded human brain tissue using CSB-PA025484HA01HU at dilution of 1:100