






## VASN Antibody

<b>Product Code</b>	CSB-PA025796OA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q6EMK4
Immunogen	Peptide sequence from Human Vasorin protein (553-572AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, WB, IHC, IF; Recommended dilution: WB:1:1000-1:5000, IHC:1:500-1:1000, IF:1:200-1:500
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
<b>Purification Method</b>	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	Vasorin (Protein slit-like 2), VASN, SLITL2
Immunogen Species	Homo sapiens (Human)
Research Area	Signal Transduction
Target Names	VASN

**Image** 



Western Blot

Positive WB detected in: HepG2 whole cell lysate, SH-SY5Y whole cell lysate, K562 whole cell lysate, 293 whole cell lysate, Hela whole cell

lysate, HL60 whole cell lysate All lanes: VASN antibody at 1:2000


Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 72 kDa Observed band size: 72 kDa















IHC image of CSB-PA025796OA01HU diluted at 1:640 and staining in paraffin-embedded human colon cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunofluorescence staining of MCF-7 cells with CSB-PA025796OA01HU at 1:213, counterstained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).