





# MSH6 Antibody

|                            |                                                                |
|----------------------------|----------------------------------------------------------------|
| <b>Product Code</b>        | CSB-PA051370                                                   |
| <b>Storage</b>             | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.  |
| <b>Uniprot No.</b>         | P52701                                                         |
| <b>Immunogen</b>           | Fusion protein of Human MSH6                                   |
| <b>Raised In</b>           | Rabbit                                                         |
| <b>Species Reactivity</b>  | Human,Mouse                                                    |
| <b>Tested Applications</b> | ELISA,WB,IHC;ELISA:1:1000-1:2000,WB:1:200-1:1000,IHC:1:10-1:50 |
| <b>Form</b>                | Liquid                                                         |
| <b>Conjugate</b>           | Non-conjugated                                                 |
| <b>Storage Buffer</b>      | -20°C, pH7.4 PBS, 0.05% NaN <sub>3</sub> , 40% Glycerol        |
| <b>Purification Method</b> | Antigen affinity purification                                  |
| <b>Isotype</b>             | IgG                                                            |
| <b>Alias</b>               | mutS homolog 6 (E. coli)                                       |
| <b>Immunogen Species</b>   | Homo sapiens (Human)                                           |
| <b>Target Names</b>        | MSH6                                                           |

## Image



The image on the left is immunohistochemistry of paraffin-embedded Human tonsil tissue using CSB-PA051370(MSH6 Antibody) at dilution 1/10, on the right is treated with fusion protein. (Original magnification: ×200)



Gel: 8%SDS-PAGE, Lysate: 40 µg, Lane 1-2: HeLa cells, 293T cells, Primary antibody: CSB-PA051370(MSH6 Antibody) at dilution 1/300, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 45 seconds