







APOE Antibody

| Product Code | CSB-PA07599A0Rb |
|---------------------|----------------------------------------------------------------------------------|
| Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Uniprot No. | P02649 |
| Immunogen | Recombinant Human Apolipoprotein E protein (21-110AA) |
| Raised In | Rabbit |
| Species Reactivity | Human |
| Tested Applications | ELISA, IHC, IF; Recommended dilution: IHC:1:1000-1:2000, IF:1:200-1:500 |
| Form | Liquid |
| Conjugate | Non-conjugated |
| Storage Buffer | Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 |
| Purification Method | >95%, Protein G purified |
| Isotype | IgG |
| Clonality | Polyclonal |
| Alias | Apolipoprotein E (Apo-E), APOE |
| Immunogen Species | Homo sapiens (Human) |
| Research Area | Neuroscience |
| Target Names | APOE |
| Image | Immunofly programme staining of Llan C2 calls |



Immunofluorescence staining of HepG2 cells with CSB-PA07599A0Rb at 1:333, counterstained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).



IHC image of CSB-PA07599A0Rb diluted at 1:1000 and staining in paraffin-embedded human brain tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.