
**Image** 





## **DKK2** Antibody

<b>Product Code</b>	CSB-PA08279A0Rb
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q9UBU2
Immunogen	Recombinant Human Dickkopf-related protein 2 protein (34-259AA)
Raised In	Rabbit
Species Reactivity	Human
<b>Tested Applications</b>	ELISA, WB, IHC; Recommended dilution: WB:1:500-1:5000, IHC:1:500-1:1000
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Storage Buffer  Purification Method	
	Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Purification Method	Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified
Purification Method	Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified  IgG
Purification Method Isotype Clonality	Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified  IgG  Polyclonal
Purification Method Isotype Clonality Alias	Constituents: 50% Glycerol, 0.01M PBS, PH 7.4  >95%, Protein G purified  IgG  Polyclonal  Dickkopf-related protein 2 (Dickkopf-2) (Dkk-2) (hDkk-2), DKK2
Purification Method Isotype Clonality Alias Immunogen Species	Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 >95%, Protein G purified  IgG  Polyclonal  Dickkopf-related protein 2 (Dickkopf-2) (Dkk-2) (hDkk-2), DKK2  Homo sapiens (Human)



Positive WB detected in: MCF-7 whole cell lysate

All lanes: DKK2 antibody at 3.9µg/ml

Secondary


Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 29 kDa Observed band size: 32 kDa










IHC image of CSB-PA08279A0Rb diluted at 1:500 and staining in paraffin-embedded human skeletal muscle tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC image of CSB-PA08279A0Rb diluted at 1:500 and staining in paraffin-embedded human lung tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.