





# NDUFA13 Antibody

|                            |                                                                  |
|----------------------------|------------------------------------------------------------------|
| <b>Product Code</b>        | CSB-PA161916                                                     |
| <b>Storage</b>             | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.    |
| <b>Uniprot No.</b>         | Q9P0J0                                                           |
| <b>Immunogen</b>           | Fusion protein of Human NDUFA13                                  |
| <b>Raised In</b>           | Rabbit                                                           |
| <b>Species Reactivity</b>  | Human,Mouse                                                      |
| <b>Tested Applications</b> | ELISA,WB,IHC;ELISA:1:2000-1:5000,WB:1:500-1:2000,IHC:1:100-1:300 |
| <b>Form</b>                | Liquid                                                           |
| <b>Conjugate</b>           | Non-conjugated                                                   |
| <b>Storage Buffer</b>      | -20°C, pH7.4 PBS, 0.05% NaN <sub>3</sub> , 40% Glycerol          |
| <b>Purification Method</b> | Antigen affinity purification                                    |
| <b>Isotype</b>             | IgG                                                              |
| <b>Immunogen Species</b>   | Homo sapiens (Human)                                             |
| <b>Target Names</b>        | NDUFA13                                                          |


## Image



The image on the left is immunohistochemistry of paraffin-embedded Human ovarian cancer tissue using CSB-PA161916(NDUFA13 Antibody) at dilution 1/50, on the right is treated with fusion protein. (Original magnification: ×200)



The image on the left is immunohistochemistry of paraffin-embedded Human colon cancer tissue using CSB-PA161916(NDUFA13 Antibody) at dilution 1/50, on the right is treated with fusion protein. (Original magnification: ×200)



Gel: 10%SDS-PAGE, Lysate: 40  $\mu$ g, Lane 1-2:  
 Mouse skeletal muscle, human hepatocellular  
 carcinoma tissue, Primary antibody: CSB-  
 PA161916(NDUFA13 Antibody) at dilution 1/350,  
 Secondary antibody: Goat anti rabbit IgG at  
 1/8000 dilution, Exposure time: 5 minutes