







## MYL3 Antibody

| <b>Product Code</b>                        | CSB-PA249474   |
|--|--|
| Storage                                    | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.                  |
| Uniprot No.                                | P08590   |
| Immunogen                                  | Fusion protein of Human MYL3   |
| Raised In                                  | Rabbit   |
| Species Reactivity                         | Human, Mouse, Rat  |
| <b>Tested Applications</b>                 | ELISA,WB,IHC;ELISA:1:2000-1:10000,WB:1:1000-1:5000,IHC:1:50-1:200              |
| Form                                       | Liquid   |
|  |  |
| Conjugate                                  | Non-conjugated   |
| Conjugate Storage Buffer                   | Non-conjugated -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol                      |
|  |  |
| Storage Buffer                             | -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol                                     |
| Storage Buffer Purification Method         | -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol  Antigen affinity purification      |
| Storage Buffer Purification Method Isotype | -20°C, pH7.4 PBS, 0.05% NaN3, 40% Glycerol  Antigen affinity purification  IgG |





The image on the left is immunohistochemistry of paraffin-embedded Human lung cancer tissue using CSB-PA249474(MYL3 Antibody) at dilution 1/45, on the right is treated with fusion protein. (Original magnification: ×200)



The image on the left is immunohistochemistry of paraffin-embedded Human liver cancer tissue using CSB-PA249474(MYL3 Antibody) at dilution 1/45, on the right is treated with fusion protein. (Original magnification: x200)




## **CUSABIO TECHNOLOGY LLC**

🕜 Tel: +1-301-363-4651 🛛 Email: cusabio@cusabio.com 🕒 Website: www.cusabio.com 💣







Gel: 10%SDS-PAGE, Lysate: 40 μg, Lane: Mouse heart tissue, Primary antibody: CSB-PA249474(MYL3 Antibody) at dilution 1/650, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 10 seconds