





# CA3 Antibody

|                            |                                                                |
|----------------------------|----------------------------------------------------------------|
| <b>Product Code</b>        | CSB-PA298167                                                   |
| <b>Storage</b>             | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.  |
| <b>Uniprot No.</b>         | P07451                                                         |
| <b>Immunogen</b>           | Fusion protein of Human CA3                                    |
| <b>Raised In</b>           | Rabbit                                                         |
| <b>Species Reactivity</b>  | Human,Mouse,Rat                                                |
| <b>Tested Applications</b> | ELISA,WB,IHC;ELISA:1:1000-1:5000,WB:1:200-1:1000,IHC:1:10-1:50 |
| <b>Form</b>                | Liquid                                                         |
| <b>Conjugate</b>           | Non-conjugated                                                 |
| <b>Storage Buffer</b>      | -20°C, pH7.4 PBS, 0.05% NaN <sub>3</sub> , 40% Glycerol        |
| <b>Purification Method</b> | Antigen affinity purification                                  |
| <b>Isotype</b>             | IgG                                                            |
| <b>Alias</b>               | Carbonic anhydrase 3                                           |
| <b>Immunogen Species</b>   | Homo sapiens (Human)                                           |
| <b>Target Names</b>        | CA3                                                            |

## Image



The image on the left is immunohistochemistry of paraffin-embedded Human thyroid cancer tissue using CSB-PA298167(CA3 Antibody) at dilution 1/25, on the right is treated with fusion protein. (Original magnification: ×200)



Gel: 10%SDS-PAGE, Lysate: 30 µg, Lane: Human fetal kidney tissue, Primary antibody: CSB-PA298167(CA3 Antibody) at dilution 1/250, Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution, Exposure time: 5 seconds