

**Image** 








## **ARHGAP32** Antibody

<b>Product Code</b>	CSB-PA415667LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	A7KAX9
Immunogen	Recombinant Human Rho GTPase-activating protein 32 protein (130-205AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, IF; Recommended dilution: IF:1:20-1:200
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
<b>Purification Method</b>	>95%, Protein G purified
Purification Method Isotype	>95%, Protein G purified  IgG
	· '
Isotype	IgG
Isotype Clonality	Polyclonal  Rho GTPase-activating protein 32, Brain-specific Rho GTPase-activating protein, GAB-associated Cdc42/Rac GTPase-activating protein, GC-GAP, GTPase regulator interacting with TrkA, Rho-type GTPase-activating protein 32, Rho/Cdc42/Rac GTPase-activating protein RICS, RhoGAP involved in the beta-catenin-N-cadherin and NMDA receptor signaling, p200RhoGAP, p250GAP,
Isotype Clonality Alias	IgG  Polyclonal  Rho GTPase-activating protein 32, Brain-specific Rho GTPase-activating protein, GAB-associated Cdc42/Rac GTPase-activating protein, GC-GAP, GTPase regulator interacting with TrkA, Rho-type GTPase-activating protein 32, Rho/Cdc42/Rac GTPase-activating protein RICS, RhoGAP involved in the beta-catenin-N-cadherin and NMDA receptor signaling, p200RhoGAP, p250GAP, ARHGAP32, GRIT, KIAA0712, RICS
Isotype Clonality Alias Immunogen Species	Polyclonal  Rho GTPase-activating protein 32, Brain-specific Rho GTPase-activating protein, GAB-associated Cdc42/Rac GTPase-activating protein, GC-GAP, GTPase regulator interacting with TrkA, Rho-type GTPase-activating protein 32, Rho/Cdc42/Rac GTPase-activating protein RICS, RhoGAP involved in the beta-catenin-N-cadherin and NMDA receptor signaling, p200RhoGAP, p250GAP, ARHGAP32, GRIT, KIAA0712, RICS  Homo sapiens (Human)



Immunofluorescence staining of SH-SY5Y cells with CSB-PA415667LA01HU at 1:33, counterstained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).