




THOC5 Antibody

Product Code	CSB-PA613425LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q13769
Immunogen	Recombinant Human THO complex subunit 5 homolog protein (2-53AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, IHC, IF; Recommended dilution: IHC:1:200-1:500, IF:1:50-1:200
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Purification Method	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	THO complex subunit 5 homolog (Functional spliceosome-associated protein 79) (fSAP79) (NF2/meningioma region protein pK1.3) (Placental protein 39.2) (PP39.2) (hTREX90), THOC5, C22orf19 KIAA0983
Immunogen Species	Homo sapiens (Human)
Research Area	Immunology
Target Names	THOC5
Image	THC image of CSR-PA613425I A01HI I diluted at



IHC image of CSB-PA613425LA01HU diluted at 1:300 and staining in paraffin-embedded human spleen tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.




CUSABIO TECHNOLOGY LLC









Immunofluorescence staining of HepG2 cells with CSB-PA613425LA01HU at 1:100, counterstained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).