


Image





CYP2A13 Antibody

| Product Code | CSB-PA621969LA01HU |
|--|---|
| Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Uniprot No. | Q16696 |
| Immunogen | Recombinant Human Cytochrome P450 2A13 protein (252-354aa) |
| Raised In | Rabbit |
| Species Reactivity | Human, Mouse |
| Tested Applications | ELISA, WB, IHC; Recommended dilution: WB:1:1000-1:5000, IHC:1:100-1:500 |
| Form | Liquid |
| Conjugate | Non-conjugated |
| | |
| Storage Buffer | Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 |
| Storage Buffer Purification Method | |
| | Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 |
| Purification Method | Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 >95%, Protein G purified |
| Purification Method | Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 >95%, Protein G purified IgG |
| Purification Method Isotype Clonality | Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 >95%, Protein G purified IgG Polyclonal |
| Purification Method Isotype Clonality Product Type | Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 >95%, Protein G purified IgG Polyclonal Polyclonal |
| Purification Method Isotype Clonality Product Type Immunogen Species | Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 >95%, Protein G purified IgG Polyclonal Polyclonal Homo sapiens (Human) |



Western Blot


Positive WB detected in: A549 whole cell lysate,

Mouse liver tissue

All lanes: CYP2A13 antibody at 1:2000

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 57 kDa Observed band size: 57 kDa



IHC image of CSB-PA621969LA01HU diluted at 1:200 and staining in paraffin-embedded human breast cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat



CUSABIO TECHNOLOGY LLC





anti-rabbit polymer IgG labeled by HRP and visualized using 0.05% DAB.