






## **APCDD1** Antibody

<b>Product Code</b>	CSB-PA814246LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q8J025
Immunogen	Recombinant Human Protein APCDD1 protein (22-288AA)
Raised In	Rabbit
Species Reactivity	Human
<b>Tested Applications</b>	ELISA, WB, IHC; Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Purification Method	Antigen Affinity Purified
Isotype	IgG
Clonality	Polyclonal
Alias	Protein APCDD1 (Adenomatosis polyposis coli down-regulated 1 protein), APCDD1, DRAPC1
Immunogen Species	Homo sapiens (Human)
Research Area	Stem Cells
Target Names	APCDD1
_	





Western Blot

Positive WB detected in: 293T whole cell lysate, MCF-7 whole cell lysate, 293 whole cell lysate, HepG2 whole cell lysate, PC-3 whole cell lysate, Hela whole cell lysate, K562 whole cell lysate

All lanes: APCDD1 antibody at 1:1000


Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 59 kDa Observed band size: 59 kDa

















IHC image of CSB-PA814246LA01HU diluted at 1:100 and staining in paraffin-embedded human heart tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC image of CSB-PA814246LA01HU diluted at 1:100 and staining in paraffin-embedded human pancreatic tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.