




CD99L2 Antibody

Product Code	CSB-PA840569LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q8TCZ2
Immunogen	Recombinant Human CD99 antigen-like protein 2 protein (26-145AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, WB, IHC, IF; Recommended dilution: WB:1:500-1:5000, IHC:1:200-1:500, IF:1:50-1:200
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Storage Buffer Purification Method	
	Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Purification Method	Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 >95%, Protein G purified
Purification Method	Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 >95%, Protein G purified IgG
Purification Method Isotype Clonality	Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 >95%, Protein G purified IgG Polyclonal CD99 antigen-like protein 2 (MIC2-like protein 1) (CD antigen CD99), CD99L2,
Purification Method Isotype Clonality Alias	Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 >95%, Protein G purified IgG Polyclonal CD99 antigen-like protein 2 (MIC2-like protein 1) (CD antigen CD99), CD99L2, MIC2L1
Purification Method Isotype Clonality Alias Immunogen Species	Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 >95%, Protein G purified IgG Polyclonal CD99 antigen-like protein 2 (MIC2-like protein 1) (CD antigen CD99), CD99L2, MIC2L1 Homo sapiens (Human)





Positive WB detected in: SH-SY5Y whole cell

All lanes: CD99L2 antibody at 4µg/ml

Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 28, 23, 21, 16, 29 kDa

Observed band size: 28 kDa















IHC image of CSB-PA840569LA01HU diluted at 1:400 and staining in paraffin-embedded human appendix tissue performed on a Leica BondTM system.



Immunofluorescence staining of A549 cells with CSB-PA840569LA01HU at 1:133, counterstained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).