








WFIKKN1 Antibody

| Product Code | CSB-PA842729OA01HU |
|---------------------|---|
| Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Uniprot No. | Q96NZ8 |
| Immunogen | Peptide sequence from Human WAP, Kazal, immunoglobulin, Kunitz and NTR domain-containing protein 1 protein (187-207AA) |
| Raised In | Rabbit |
| Species Reactivity | Human |
| Tested Applications | ELISA, IHC; Recommended dilution: IHC:1:200-1:500 |
| Form | Liquid |
| Conjugate | Non-conjugated |
| Storage Buffer | Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 |
| Purification Method | >95%, Protein G purified |
| Isotype | IgG |
| Clonality | Polyclonal |
| Alias | WAP, Kazal, immunoglobulin, Kunitz and NTR domain-containing protein 1 (Growth and differentiation factor-associated serum protein 2) (GASP-2) (hGASP-2) (WAP, follistatin, immunoglobulin, Kunitz and NTR domain-containing protein), WFIKKN1, C16orf12 GASP2 WFIKKN |
| Immunogen Species | Homo sapiens (Human) |
| Research Area | Cell Biology |
| Target Names | WFIKKN1 |
| Image | IHC image of CSR-PA8/27290A01HI I diluted at |



IHC image of CSB-PA842729OA01HU diluted at 1:300 and staining in paraffin-embedded human liver cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.