




# NEDD4L Antibody

<b>Product Code</b>	CSB-PA850411LA01HU
<b>Storage</b>	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
<b>Uniprot No.</b>	Q96PU5
<b>Immunogen</b>	Recombinant Human E3 ubiquitin-protein ligase NEDD4-like protein (289-494AA)
<b>Raised In</b>	Rabbit
<b>Species Reactivity</b>	Human
<b>Tested Applications</b>	ELISA, WB, IF; Recommended dilution: WB:1:500-1:5000, IF:1:50-1:200
<b>Form</b>	Liquid
<b>Conjugate</b>	Non-conjugated
<b>Storage Buffer</b>	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
<b>Purification Method</b>	>95%, Protein G purified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Alias</b>	E3 ubiquitin-protein ligase NEDD4-like (EC 2.3.2.26) (HECT-type E3 ubiquitin transferase NED4L) (NEDD4.2) (Nedd4-2), NEDD4L, KIAA0439 NEDL3
<b>Immunogen Species</b>	Homo sapiens (Human)
<b>Research Area</b>	Epigenetics and Nuclear Signaling
<b>Target Names</b>	NEDD4L

## Image



### Western Blot

Positive WB detected in: 293 whole cell lysate,

Jurkat whole cell lysate, HeLa whole cell lysate

All lanes: NEDD4L antibody at 3µg/ml


### Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 112, 105, 101, 99, 111,

110, 97 kDa

Observed band size: 112 kDa



Immunofluorescence staining of HepG2 cells with CSB-PA850411LA01HU at 1:100, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).