ARHGAP18 Antibody

|	
Product Code	CSB-PA850791LA01HU
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q8N392
Immunogen	Recombinant Human Rho GTPase-activating protein 18 protein (389-577AA)
Raised In	Rabbit
Species Reactivity	Human, Rat, Mouse
Tested Applications	ELISA, WB, IHC, IF; Recommended dilution: WB:1:500-1:5000, IHC:1:200-1:500, IF:1:50-1:200
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Purification Method	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	Rho GTPase-activating protein 18 (MacGAP) (Rho-type GTPase-activating protein 18), ARHGAP18
Immunogen Species	Homo sapiens (Human)
Research Area	Signal Transduction
Target Names	ARHGAP18
Image	

Western Blot
Positive WB detected in: Hela whole cell lysate, NIH/3T3 whole cell lysate, A549 whole cell lysate, Rat lung tissue, Rat kidney tissue, Mouse heart tissue
All lanes: ARHGAP18 antibody at 4.5μg/ml
Secondary
Goat polyclonal to rabbit IgG at 1/50000 dilution
Predicted band size: 75, 71 kDa
Observed band size: 75 kDa

IHC image of CSB-PA850791LA01HU diluted at 1:200 and staining in paraffin-embedded human small intestine tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

Immunofluorescence staining of HepG2 cells with CSB-PA850791LA01HU at 1:66, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).