






CELSR3 Antibody

| Product Code | CSB-PA852817LA01HU |
|---------------------|--|
| Storage | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Uniprot No. | Q9NYQ7 |
| Immunogen | Recombinant Human Cadherin EGF LAG seven-pass G-type receptor 3 protein (514-718AA) |
| Raised In | Rabbit |
| Species Reactivity | Human |
| Tested Applications | ELISA, IHC; Recommended dilution: IHC:1:200-1:500 |
| Form | Liquid |
| Conjugate | Non-conjugated |
| Storage Buffer | Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4 |
| Purification Method | >95%, Protein G purified |
| Isotype | IgG |
| Clonality | Polyclonal |
| Alias | Cadherin EGF LAG seven-pass G-type receptor 3 (Cadherin family member 11) (Epidermal growth factor-like protein 1) (EGF-like protein 1) (Flamingo homolog 1) (hFmi1) (Multiple epidermal growth factor-like domains protein 2) (Multiple EGF-like domains protein 2), CELSR3, CDHF11 EGFL1 FMI1 KIAA0812 MEGF2 |
| Immunogen Species | Homo sapiens (Human) |
| Research Area | Signal Transduction |
| Target Names | CELSR3 |
| Image | IHC image of CSR-PA852817I A01HI I diluted at |



IHC image of CSB-PA852817LA01HU diluted at 1:300 and staining in paraffin-embedded human liver cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.