




# XAB2 Antibody

|                            |                                                                                                        |
|----------------------------|--------------------------------------------------------------------------------------------------------|
| <b>Product Code</b>        | CSB-PA862073LA01HU                                                                                     |
| <b>Storage</b>             | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.                                          |
| <b>Uniprot No.</b>         | Q9HCS7                                                                                                 |
| <b>Immunogen</b>           | Recombinant Human Pre-mRNA-splicing factor SYF1 protein (750-840AA)                                    |
| <b>Raised In</b>           | Rabbit                                                                                                 |
| <b>Species Reactivity</b>  | Human                                                                                                  |
| <b>Tested Applications</b> | ELISA, WB, IHC, IF; Recommended dilution: WB:1:500-1:5000, IHC:1:200-1:500, IF:1:50-1:200              |
| <b>Form</b>                | Liquid                                                                                                 |
| <b>Conjugate</b>           | Non-conjugated                                                                                         |
| <b>Storage Buffer</b>      | Preservative: 0.03% Proclin 300<br>Constituents: 50% Glycerol, 0.01M PBS, pH 7.4                       |
| <b>Purification Method</b> | >95%, Protein G purified                                                                               |
| <b>Isotype</b>             | IgG                                                                                                    |
| <b>Clonality</b>           | Polyclonal                                                                                             |
| <b>Alias</b>               | Pre-mRNA-splicing factor SYF1, Protein HCNP, XPA-binding protein 2, XAB2, HCNP, KIAA1177, SYF1, PP3898 |
| <b>Immunogen Species</b>   | Homo sapiens (Human)                                                                                   |
| <b>Research Area</b>       | Epigenetics and Nuclear Signaling                                                                      |
| <b>Target Names</b>        | XAB2                                                                                                   |

## Image



### Western Blot

Positive WB detected in: HeLa whole cell lysate, K562 whole cell lysate, A549 whole cell lysate


All lanes: XAB2 antibody at 3µg/ml

### Secondary


Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 101 kDa


Observed band size: 101 kDa



IHC image of CSB-PA862073LA01HU diluted at 1:400 and staining in paraffin-embedded human brain tissue performed on a Leica Bond<sup>TM</sup> system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC image of CSB-PA862073LA01HU diluted at 1:400 and staining in paraffin-embedded human colon cancer performed on a Leica Bond<sup>TM</sup> system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunofluorescence staining of PC-3 cells with CSB-PA862073LA01HU at 1:133, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).